

Universal lattices and unbounded rank expanders

Martin Kassabov

Abstract

We study the representations of non-commutative universal lattices and use them to compute lower bounds for the τ -constant for the commutative universal lattices $G_{d,k} = \mathrm{SL}_d(\mathbb{Z}[x_1, \dots, x_k])$ with respect to several generating sets.

As an application of the above result we show that the Cayley graphs of the finite groups $\mathrm{SL}_{3k}(\mathbb{F}_p)$ can be made expanders using suitable choice of the generators. This provides the first examples of expander families of groups of Lie type where the rank is not bounded and gives a natural (and explicit) counter examples to two conjectures of Alex Lubotzky and Benjamin Weiss.

Introduction

The groups $\mathrm{SL}_d(\mathcal{O})$, where $\mathcal{O}$ is a ring of integers in a number field K , have many common — they all have Kazhdan property T , have a finite congruence kernel and super rigidity. In [14], Y. Shalom conjectured that many of these properties are inherited from the group $\mathrm{SL}_d(\mathbb{Z}[x])$. He called the groups $\mathrm{SL}_d(\mathbb{Z}[x_1, \dots, x_k])$ *universal lattices*. Almost nothing is known about the representation theory of these groups. It is conjectured that the universal lattices have property T , it was known that these groups have property τ , see [8].

In this paper we study the noncommutative analogs of these groups, i.e., the subgroups of $\mathrm{GL}_d(\mathbb{Z}\langle x_1, \dots, x_k \rangle)$ generated by all elementary matrices with coefficients in the free associative ring on k generators $\mathbb{Z}\langle x_1, \dots, x_k \rangle$. Using that these groups can be mapped onto many universal lattices we obtain new bounds for the τ -constant for the groups $\mathrm{SL}_d(\mathbb{Z}[x_1, \dots, x_k])$ with respect to several generating sets.

Let G be a topological group and consider the space $\tilde{G}$ of all equivalence classes of unitary representations of G on some Hilbert space $\mathcal{H}$. This space has a naturally defined topology, called the Fell topology, as explained in [13] §1.1 or [11] Chapter 3 for example. Let 1_G denote the trivial 1-dimensional

2000 Mathematics Subject Classification: Primary 20F69; Secondary 05C25, 05E15, 19C20, 20C33, 20G35, 20H05, 22E40, 22E55.

Key words and phrases: universal lattices, property T , property τ , Kazhdan constant, τ -constant, expanders, unbounded rank expanders, Lubotzky-Weiss conjectures.

representation of G and let $\tilde{G}_0$ be the set of representations in $\tilde{G}$ which do not contain 1_G as a subrepresentation (i.e., do not have invariant vectors).

Definition 1 *A group G is said to have Kazhdan property T if 1_G is isolated from $\tilde{G}_0$ in the Fell topology of $\tilde{G}$.*

A discrete group with property T is finitely generated: see [10]. Since in this paper we shall be concerned mostly with discrete groups we give the following equivalent reformulation of property T :

Equivalent definition: Let G be a discrete group generated by a finite set S . Then G has the Kazhdan property T if there is $\epsilon > 0$ such that for every nontrivial irreducible unitary representation $\rho : G \rightarrow U(\mathcal{H})$ on a Hilbert space $\mathcal{H}$ and every vector $v \neq 0$ there is some $s \in S$ such that $\|\rho(s)v - v\| > \epsilon\|v\|$. The largest ϵ with this property is called the *Kazhdan constant* for S and is denoted by $\mathcal{K}(G; S)$.

The property T depends only on the group G and does not depend on the choice of the generating set S , however the Kazhdan constant depends also on the generating set.

Property T implies certain group theoretic conditions on G (finite generation, FP, FAB etc) and can be used for construction of *expanders* from the finite images of G . For this last application the following weaker property τ is sufficient:

Let $\tilde{G}^f$ and $\tilde{G}_0^f$ denote the finite representations of $\tilde{G}$, resp. $\tilde{G}_0$ (i.e., the representations which factor through a finite index subgroup).

Definition 2 *A group G is said to have property τ if 1_G is isolated from $\tilde{G}_0^f$ in the induced Fell topology of $\tilde{G}^f$. Equivalently: the group G with the profinite topology and its profinite completion $\hat{G}$ has property T.*

Again, for a discrete finitely generated¹ group G there is an equivalent definition of these properties:

Equivalent definition: Let G be an discrete group generated by a set S . Then G has the property τ if there is $\epsilon > 0$ such that for every nontrivial finite irreducible unitary representation $\rho : G \rightarrow U(\mathcal{H})$ on a Hilbert space $\mathcal{H}$ and every vector $v \neq 0$ there is some $s \in S$ such that $\|\rho(s)v - v\| > \epsilon\|v\|$. The largest ϵ with this property is called the τ -constant and is denoted by $\tau(G; S)$.

Property τ is not interesting for groups which do not have many finite quotients. All the groups we are going to work with are residually finite and have enough finite factors.

As mentioned above, properties T and τ can be used for the construction of families of *expanding graphs*. We need more terminology.

¹A discrete group with property τ may not be finitely generated. However, all the groups we are going to work with are f.g., and there is no need to define τ -constant for not finitely generated groups.

A finite graph Γ is called an ϵ -expander for some $\epsilon \in (0, 1)$ if for any subset $A \subseteq \Gamma$ of size at most $|\Gamma|/2$ we have $|\partial(A)| > \epsilon|A|$ (where $\partial(A)$ is the set of vertices of $\Gamma \setminus A$ of edge distance 1 to A). The largest such ϵ is called the expanding constant of Γ . Constructing families of ϵ -expanders with large expanding constant ϵ and bounded valency is an important practical problem in computer science. For an excellent introduction to the subject we refer the reader to the book [11] by Lubotzky. The following equivalent definition of property τ illustrates one of the few known approaches to explicit construction of expanders:

Proposition 3 ([11], **Theorem 4.3.2**) *Let G be a finitely generated discrete group. Then G has property τ iff for any set of generators S there exists an $\epsilon = \epsilon(S) > 0$ such that the Cayley graphs $\mathcal{C}(G_i, S_i)$ of the finite images of G_i of G (with respect to the images S_i of S) form a family of ϵ -expanders. The largest $\epsilon_0(S)$ with this property is related to the τ -constant $\tau(G, S)$ defined above. In particular we have $\epsilon_0(S) \geq \tau(G, S)^2/4$.*

Our approach to property τ is inspired by a paper by Shalom [14] which relates property T to *bounded generation property* of high rank Chevalley groups over rings of integers. In this paper we will work only with the group $\text{EL}_d(R)$ for $d \geq 3$ which correspond to the Dynkin diagram A_{d-1} . Similar arguments also work for high rank the Chevalley groups which arise from others Dynkin diagrams.

Let R be an associative ring with unit, and for $i \neq j \in \{1, 2, \dots, d\}$ let $E_{i,j}$ denote the set of elementary $d \times d$ matrices $\{\text{Id} + r \cdot e_{i,j} \mid r \in R\}$. Set $E = E(R) = \bigcup_{i \neq j} E_{i,j}$ and let $\text{EL}(d; R) = \text{EL}_d(R)$ be the subgroup of the multiplicative group of the ring of $d \times d$ matrices over R generated by $E(R)$.

Definition 4 *The group $G = \text{EL}_d(R)$ is said to have bounded elementary generation property if there is a number $N = N(R)$ such that every element of G can be written as a product of at most N elements from the set E .*

Examples of infinite rings R satisfying the above definition are rings of integers $\mathcal{O}$ in number fields K (for $d \geq 3$), see [3]. In this classical case this property is known as bounded generation because each group $E_{i,j} \simeq (\mathcal{O}, +)$ is a product of finitely many cyclic groups.

The following theorem is a generalization of a result of Y. Shalom (see Theorem 3.4 in [14]) to non-commutative rings:

Theorem 5 *Suppose that $d \geq 3$ and R is a finitely generated associative ring such that $\text{EL}_d(R)$ has bounded elementary generation property. Then $\text{EL}_d(R)$ has property T (as a discrete group).*

Moreover there is an explicit lower bound for the Kazhdan constant $\mathcal{K}(G, S)$ with respect to a specific generating set S .

A very interesting conjecture is whether $\text{EL}_d(\mathbb{Z}[x]) = \text{SL}_d(\mathbb{Z}[x])$ and other universal lattices, has bounded elementary generation property. This together

with Theorem 5 implies that the universal lattices has property T . In [8] it was shown that the groups

$$G_{d,k} := \text{EL}_d(\mathbb{Z}[x_1, \dots, x_k]) = \text{SL}_d(\mathbb{Z}[x_1, \dots, x_k])$$

have property τ . We believe that the methods in this paper can be used to show that the universal lattice in the non-commutative case have property τ . In order to show that we need several results about the structure of finite non-commutative ring R and the K -groups $K_1(R)$ and $K_2(R)$.

Theorem 5 has several applications:

Tau Constant for $\text{SL}_d(\mathbb{Z}[x_1, \dots, x_k])$.

In [8], the commutative universal lattices are shown to have property τ and the τ -constant for $G_{d,k} = \text{SL}_d(\mathbb{Z}[x_1, \dots, x_k])$ is estimated. Theorem 5 from [8] gives a lower bound for the τ -constant with respect to the generating set $S_{d,k}$, which is asymptotically $O(d^{-2}22^{-k})$.

However, it is possible to improve this estimate to $O(d^{-1/2}(1 + k/d)^{-3/2})$. We state this result as Theorem 6, its proof is an extension of the one in [8] and combines the ideas from [7], namely relative property T of certain groups and the use of *generalized elementary matrices*, see Theorem 2.9 and Lemma 3.2.

Theorem 6 *The commutative universal lattices*

$$G_{d,k} = \text{EL}_d(\mathbb{Z}[x_1, \dots, x_k]) = \text{SL}_d(\mathbb{Z}[x_1, \dots, x_k])$$

have property τ . The τ -constant with respect to the generating set $S_{d,k}$ consisting of all elementary matrices with ± 1 of the diagonal and the ones with $\pm x_i$ next to the main diagonal, satisfies

$$\tau(G_{d,k}, S_{d,k}) \geq \frac{1}{800\sqrt{d}(1 + (k/d)^{3/2})}.$$

It is interesting if this bound is asymptotically exact. Using a natural representation of $\text{SL}_d(\mathbb{Z}[x_1, \dots, x_k])$ in the Hilbert space $l^2(\mathbb{F}_2^d)$ it can be seen that

$$\tau(G_{d,k}, S_{d,k}) \leq \sqrt{\frac{2}{d}},$$

which shows that if k is fixed then the bound for τ -constant in Theorem 6 is asymptotically exact. We are not aware of any nontrivial upper bound for $\tau(G_{d,k}, S_{d,k})$ which depends on the number of generators k .

If one modifies the generating set $S_{d,k}$ then the τ -constant can be improved, in fact Theorem 11 gives that for many d there exist a finite generating set $S'_{d,k}$ such that $\tau(G_{d,k}, S'_{d,k})$ depends only on k/d .

Unbounded rank expanders.

Theorem 5 has another interesting application. In [12] Lubotzky and Weiss made the following conjectures:

Conjecture 7 ([12], Problem 5.1) *Let Γ_i be a family of finite groups and let Σ_i and Σ'_i be two generating sets of the group Γ_i of uniformly bounded size. Suppose that the Cayley graphs $\mathcal{C}(\Gamma_i, \Sigma_i)$ form an expander family. Is it true that $\mathcal{C}(\Gamma_i, \Sigma'_i)$ is also an expander family?*

Conjecture 8 ([12], Conjecture 5.4) *Let K be a compact group. If Γ_1 and Γ_2 are two finitely generated dense subgroups of K with Γ_1 amenable and Γ_2 having property T then the group K is finite.*

In [1] Alon, Lubotzky and Wigderson showed that Conjecture 7 is false using zig-zag products. Their construction is based on probabilistic arguments and does not give explicit generating sets which make the Cayley graphs expanders. Using Theorem 5, we are able to construct more natural counter examples of this conjecture, moreover in our counter example all groups and generating sets are explicit and the expanding constants can be estimated.

Consider the family of groups $\Gamma_{n,p} = \mathrm{SL}_n(\mathbb{F}_p)$, $n > 3$. Let Σ_n be the generating set of $\Gamma_{n,p}$ consisting of

$$A_n = \sum_{i=1}^{n-1} e_{i,i+1} + (-1)^{n-1} e_{n,1} \quad \text{and} \quad B_n = \mathrm{Id}_n + e_{1,2}$$

and their inverses. In [12], it is shown that the Kazhdan constants of these groups are bounded from above by

$$\mathcal{K}(\Gamma_{n,p}, \Sigma_n) \leq \sqrt{2/n},$$

which implies that any infinite subset of the Cayley graphs $\mathcal{C}(\Gamma_{n,p}, \Sigma_n)$ for fixed p and different n , are not expanders.² In view of Conjecture 7 this implies that any infinite family of the Cayley graphs of $\Gamma_{n,p}$ are not expanders with respect to any generating set of uniformly bounded size.

Using non-commutative universal lattices $G_{g,k}^n$ it is possible to find generating sets Σ'_{3l} of $\Gamma_{3l,p}$ such that the Cayley graphs are expanders:

Let us observe that the matrix algebras $\mathrm{Mat}_l(\mathbb{F}_p)$ can be generated as rings by 2 elements, i.e., all these algebras are quotients of the ring $\mathbb{Z}\langle x_1, x_2 \rangle$. It is well known that

$$\mathrm{Mat}_d(\mathrm{Mat}_l(R)) \simeq \mathrm{Mat}_{dl}(R).$$

These two observations allow us to consider the groups $\mathrm{SL}_{3l}(\mathbb{F}_p)$ as quotients of $\mathrm{EL}_3(\mathbb{Z}\langle x_1, x_2 \rangle)$. Theorem 5 gives us that the group $\mathrm{EL}_3(\mathbb{Z}\langle x_1, x_2 \rangle)$ almost has property τ with respect to the generating set $S_{3,2}$ consisting of 28 elements. If

²It is interesting to note that although these Cayley are not expanders, they have a logarithmic diameter, see [9]

we project this generating set to a generating set Σ'_{3l} of $\mathrm{SL}_{3l}(\mathbb{F}_p)$ we obtain that Kazhdan constant is uniformly bounded from below and therefore the Cayley graphs $\mathcal{C}(\mathrm{SL}_{3l}(\mathbb{F}_p), \Sigma'_{3l})$ form an expander family.³ In fact we have shown that

Theorem 9 *For all primes p and $l \geq 1$, there exists a generating set Σ'_{3l} of the group $\mathrm{SL}_{3l}(\mathbb{F}_p)$ with 28 elements such that the Kazhdan constants*

$$\mathcal{K}(\mathrm{SL}_{3l}(\mathbb{F}_p), \Sigma'_{3l}) > 1/400 \quad \text{for all } l, p,$$

which implies that the Cayley graphs $\mathcal{C}(\mathrm{SL}_{3l}(\mathbb{F}_p), \Sigma'_{3l})$ form a family of expanders with expanding constant at least 1.5×10^{-6} .

Theorem 9 gives a positive answer to the question [13, Problem 5.27], whether the Cayley graphs of $\mathrm{SL}_n(\mathbb{F}_p)$ for fixed p can be made expanders.

Similar construction almost allows us to disprove Conjecture 8 — it is known (see [12]) that in the product $\prod_n \mathrm{SL}_n(\mathbb{F}_p)$ there is a finitely generated dense amenable subgroup. We can show that there is also a dense subgroup which has property τ .⁴ In order to simplify the argument we will construct a counter example inside the product $\prod_l \mathrm{SL}_{3l}(\mathbb{F}_p)$:

Theorem 10 *Let $p > 2$ be a prime and let $\tilde{K}^p$ be the infinite compact group given by the product:*

$$\tilde{K}^p := \prod_{l \geq 3} \mathrm{SL}_{3l}(\mathbb{F}_p)$$

with the the product topology. There exist finitely generated dense subgroups Γ_1 and Γ_2 of K such that Γ_1 is amenable and Γ_2 has property τ .

This idea can be applied to a slightly more general situation. There is a surjection $R_{s+2}^n \rightarrow \mathrm{Mat}_l(R_{sl}^n)$, which gives a surjection $G_{3,s+2}^n \rightarrow G_{3l,sl}^n$. Using this surjection we can prove that:

Theorem 11 *There exists a set $S'_{3l,sl}$ with $28+8s$ elements generating the group $G_{3l,sl} = \mathrm{SL}_{3l}(\mathbb{Z}[x_1, \dots, x_{sl}])$ such that the τ -constant $\tau(G_{3l,sl}, S'_{3l,sl})$ depends only on s .*

A few words about the structure of the rest of the paper: Section 1 contains the proof of Theorems 5 and 6 modulo Lemmas 1.1 and 1.2. In section 2 we prove some technical results about relative property T and relative Kazhdan constants. In section 3 we prove uniform bounded elementary generation of the group $\mathrm{SL}_d(\bar{R})$ for some finite $\bar{R}$. The last section 4 is devoted to applications of Theorem 5 – a construction of expander family of Lie groups of unbounded rank and a counter examples of the conjectures of A. Lubotzky and B. Wiess.

³This argument can be generalized to show that for any $n \geq 3$ and any prime p , there exists a generating set $\Sigma''_{n,p}$ of $\mathrm{SL}_n(\mathbb{F}_p)$, such that the Cayley graphs $\mathcal{C}(\mathrm{SL}_n(\mathbb{F}_p), \Sigma''_{n,p})$ for all n and p form an expander family.

⁴It is possible that this group also have property T , but we are unable to prove it.

Acknowledgements

I am grateful to R. K. Dennis, A. Lubotzky, N. Nikolov, Y. Shalom and L. Vaserstain for introducing me to the subject and for useful discussions. I additionally wish to thank Nikolay Nikolov for providing the the proof of Lemma 1.2 and others results in section 3.

Notations

Let $R_k^n := \mathbb{Z}\langle x_1, \dots, x_k \rangle$ be the free associative ring on k (non-commutative) generators, and let $R_k := R_k^n / [R_k^n, R_k^n] = \mathbb{Z}[x_1, \dots, x_k]$ be its abelinization, which is isomorphic to the polynomial ring on k variables with coefficients in $\mathbb{Z}$.

Let R be an associative ring with 1, consider the ring $\text{Mat}_d(R)$ of $d \times d$ matrices of R . With $\text{GL}_d(R)$ we will denote the multiplicative group $\text{Mat}_d(R)$, i.e., the group of invertible $d \times d$ matrices with coefficients in the ring R . We will assume that $d \geq 3$.

For $i \neq j \in \{1, 2, \dots, d\}$ let $E_{i,j}$ denote the set of elementary $d \times d$ matrices of the form $\{\text{Id} + r \cdot e_{i,j} \mid r \in R\}$, it is clear that these matrices are invertible. Also set $E = E(R) = \bigcup_{i \neq j} E_{i,j} \subset \text{GL}_d(R)$ and let $\text{EL}(d; R) = \text{EL}_d(R)$ be the subgroup in $\text{GL}_d(R)$ generated by $E(R)$. In the case of finitely generated commutative ring R this is a subgroup of $\text{SL}_d(R)$. For the rest of this paper we will denote $G_{d,k}^n := \text{EL}(d; R_k^n)$ and $G_{d,k} := \text{EL}(d; R_k)$. Unless we explicitly need to specify the number of generators of the ring R_k^n or the size of matrices $G_{d,k}^n$ and $G_{d,k}$ we will denote R_k^n by R^n , R_k by R , $G_{d,k}^n$ by G^n and $G_{d,k}$ by G .

The groups $G_{d,k}^n$ and $G_{d,k}$ are generated⁵ by the set $S_{d,k} = F_1 \cup F_2$, where F_1 is the set of $2(d^2 - d)$ elementary matrices with ± 1 off the diagonal and F_2 is the set $4k(d - 1)$ elementary matrices $\text{Id} \pm x_l e_{ij}$ with $|i - j| = 1$ and $1 \leq l \leq k$.

For any $r \in R$ and $1 \leq i \neq j \leq d$, with $e_{ij}(r)$ we will denote the matrix $\text{Id} + r \cdot e_{ij} \in \text{EL}_d(R) \subset \text{GL}_d(R)$. If r is a unit of the ring R , $e_{ii}(r)$ will denote the matrix $\text{Id} + (r - 1) \cdot e_{ii} \in \text{GL}_d(R)$.

We will call the elements in $E(R)$ elementary matrices. Any matrix in $\text{EL}_{p+q}(R)$ of the form

$$\begin{pmatrix} \text{Id}_p & * \\ 0 & \text{Id}_q \end{pmatrix} \quad \text{or} \quad \begin{pmatrix} \text{Id}_q & 0 \\ * & \text{Id}_p \end{pmatrix}$$

where 0 and $*$ are blocks of sizes $p \times q$ and $q \times p$ respectively is called a *generalized elementary matrix* (abbreviated to GEM).

We will also assume that all rings are associative with 1.

1 Proof of Theorems 5 and 6

Proof of Theorem 5: Let $\rho : \text{EL}_d(R) \rightarrow U(\mathcal{H})$ be a unitary representation of the group $\text{EL}_d(R)$ for some k -generated associative ring R , which have the

⁵The sets $S_{d,k}$ generate the groups $G_{d,k}$ and $G_{d,k}^n$ provided that $d \geq 3$. If $d = 2$ these groups are not finitely generated.

bounded elementary generation property. Suppose that $v \in \mathcal{H}$ is an ϵ -almost invariant unit vector for the set S , where $\epsilon \leq K_{d,k}$, here $K_{d,k}$ is a constant which will be determined later.

The next lemma is a corollary to Theorems 2.2 and 2.9, proved in Section 2.

Lemma 1.1 *There exist constants*

$$M(k) < 3\sqrt{2}(\sqrt{k} + 3) \quad \text{and} \quad M_d(k) < 6\sqrt{2}(\sqrt{k} + 3) + \sqrt{3d}$$

with the following properties: Let ρ be a unitary representation of the group $G_{d,k}^n = \text{EL}(d; R_k^n)$ in a Hilbert space $\mathcal{H}$. Let $v \in \mathcal{H}$ be a unit vector such that $\|\rho(g)v - v\| < \epsilon$ for $g \in S_{d,k}$. Then:

- a) We have that $\|\rho(g)v - v\| \leq 2M(k)\epsilon$ for every elementary matrix g .*
- b) We have that $\|\rho(g)v - v\| \leq 2M_d(k)\epsilon$ for every generalized elementary matrix g .*

To finish the proof of Theorem 5 we use an argument due to Y. Shalom from [14]. Let g be any element in the group $\text{EL}_d(R)$. Using the bounded elementary generation property of R , we can write

$$g = \prod_{s=1}^{BE_d(R)} u_s,$$

where u_s are elementary matrices. Finally, we can use Lemma 1.1 to obtain:

$$\|\rho(g)v - v\| \leq \sum_{s=1}^{BE_d(R)} \|\rho(u_s)v - v\| < \sum_{s=1}^{BE_d(R)} 2M(k)\epsilon = 2BE_d(R)M(k)\epsilon.$$

If we chose $K_{d,k} = (\sqrt{2}BE_d(R)M(k))^{-1}$ and use that $\epsilon \leq K_{d,k}$ we obtain

$$\|\rho(g)v - v\| < 2BE_d(R)M(k)K_{d,k} = \sqrt{2}.$$

This show that every element in the group $\text{EL}_d(R)$ moves the vector v by less than $\sqrt{2}$. Let V denote the G^n -orbit of the vector v in $\mathcal{H}$. The center of mass c_V of the set V is invariant under the action of G and is not zero because the orbit V lies entirely in the half space $\{v' \in \mathcal{H} \mid \Re(v, v') > 0\}$. Therefore $\mathcal{H}$ contains a nontrivial G -invariant vector, which completes the proof that the group $\text{EL}_d(R)$ has the property T and also provides a lower bound for the Kazhdan constant. $\square$

Proof of Theorem 6: We can not apply Theorem 5 directly because it is not known whether the group $G_{d,k} = \text{SL}_d(\mathbb{Z}[x_1, \dots, x_k])$ has bounded elementary generation property. We want only to estimate the τ -constant for this group, not the Kazhdan constant, and it is sufficient to show that all finite images of $G_{d,k}$ have uniform bounded elementary generation property.

Let $\rho : G_{d,k} \rightarrow U(\mathcal{H})$ be a unitary representation of the universal lattice $G_{d,k}$ which factors through a finite index subgroup. Suppose that $v \in \mathcal{H}$ is an ϵ -almost invariant unit vector for the set $S_{d,k}$, where $\epsilon \leq T_{d,k}$. Here $T_{d,k}$ is a constant depending on d and k which will be fixed later.

Let $H = \ker \rho < G$, this is a normal subgroup of finite index in G . Define subgroups $H_{i,j}$ of H and subsets $U_{i,j}$ of the ring R for $i \neq j$ as follows

$$H_{i,j} := H \cap E_{i,j}; \quad U_{i,j} := \{r \in R \mid 1 + r \cdot e_{i,j} \in H\}.$$

Using elements in the Weyl group we can see that the subgroups $E_{i,j}$ are pairwise conjugate in $\mathrm{SL}_d(R)$. This gives us that the sets $U_{i,j}$ do not depend on the indices i, j . Using commutation with a suitable elementary matrix we can see that $U_{i,j}$ is also closed under multiplication by elements in R , i.e., $U = U_{i,j}$ is an ideal of the ring R . This ideal is of finite index in R because H is a subgroup of finite index in G .

Let $\mathrm{EL}(d; U)$ be the normal subgroup in G generated by $H_{i,j}$:

$$\mathrm{EL}(d; U) := \langle H_{i,j} \rangle^G < H$$

Let $SK_1(R, U; d) = G(U)/\mathrm{EL}(d; U)$ where $G(U)$ is the principal congruence subgroup of G modulo U , i.e., the matrices in G congruent to Id_d modulo U . We have the following diagram

$$\begin{array}{c} 1 \rightarrow SK_1(R, U; d) \rightarrow G/\mathrm{EL}(d; U) \rightarrow G(R/U) = \mathrm{SL}_d(\bar{R}) \rightarrow 1 \\ \downarrow \\ G/H \\ \downarrow \\ 1 \end{array}$$

where the row and the column are exact.

The next result is known in the case of commutative rings (cf. [5], Remark 10 in the commutative case). Its proof in case of matrices over finite commutative ring is slightly more complicated:

Lemma 1.2 *Let $\bar{R} = \mathrm{Mat}_t(R')$ be a matrix ring over a finite commutative ring R' . The group $\mathrm{EL}_d(\bar{R}) = \mathrm{SL}_{dt}(R')$ has uniform bounded elementary generation property, i.e., number $BE_d(\bar{R})$ of elementary matrices needed to express any element in $\mathrm{EL}_d(\bar{R})$ is depends only on d and is independent of the ring $\bar{R}$.*

In fact every matrix in $\mathrm{EL}_d(\bar{R})$ can be written as a product of 38 generalized elementary matrices.

It remains to deal with the group $SK_1(R, U; d)$. It is always a finite abelian group and it measures the departure of $\mathrm{EL}(d; U)$ from being a congruence subgroup. The following lemma is proved in [8]:

Lemma 1.3 ([8], Lemma 10) *Each element of the groups $K_2(R/U)$ is a product of at most $k + 2$ Steinberg symbols $\{a, b\}$, for a, b invertible elements in R/U . The same is true for its image $SK_1(R, U; d)$.*

For a $u, u' \in R$ such that $uu' = 1 \pmod{U}$ and a pair of indices i, j consider the elements

$$w_{i,j}(u, u') := e_{i,j}(u)e_{j,i}(-u')e_{i,j}(u) \text{ and } h_{i,j}(u, u') := w_{i,j}(u, u')w_{i,j}(-1, -1),$$

in $\text{SL}_d(R)/\text{EL}(d; U)$. It is clear that the elements $w_{i,j}(u, u')$ and $h_{i,j}(u, u')$ depend only on the image of u in R/U and we can denote them $w_{i,j}(\bar{u})$ and $h_{i,j}(\bar{u})$. For any units $\bar{u}, \bar{v} \in R/U$ we define the Steinberg symbol

$$\{\bar{u}, \bar{v}\}_{i,j} = h_{i,j}(\bar{u}\bar{v})h_{i,j}(\bar{u})^{-1}h_{i,j}(\bar{v})^{-1} \in \text{SL}_d(R)/\text{EL}(d; U).$$

It can be shown that the element $\{\bar{u}, \bar{v}\}_{i,j}$ is independent on the choice of i, j and lies in the kernel of $SK_1(R, U; d)$. This element is called a Steinberg symbol and is denoted by $\{\bar{u}, \bar{v}\}$. By definition each Steinberg symbol can be written as a product of 13 elementary matrices.

For elements $u_i, u'_i \in R$, $i = 1, \dots, p \leq d/2$ such that $u_i u'_i = 1 \pmod{U}$ consider the elements

$$\tilde{w}(u_i; u'_i) = \prod_i e_{i,p+i}(u_i) \cdot \prod_i e_{p+i,i}(-u'_i) \cdot \prod_i e_{i,p+i}(u_i)$$

and

$$\tilde{h}(u_i; u'_i) := \tilde{w}(u_i; u'_i)\tilde{w}(-1, \dots; -1, \dots)$$

Using the definition of the Steinberg symbols it can be seen that

$$\prod_i \{\bar{u}_i, \bar{v}_i\} = \tilde{h}(u_i v_i; u'_i v'_i) \tilde{h}(u_i; u'_i)^{-1} \tilde{h}(v_i; v'_i)^{-1}.$$

Therefore any product of $p = \lfloor d/2 \rfloor$ Steinberg symbols can be written as a product of 13 generalized elementary matrix – here we have used that $\prod_i e_{i,p+i}(u_i)$ is a GEM. This gives us that any element in $SK_1(R, U)$ can be written as

$$13 \lceil (k+2)/p \rceil \leq 13(3 + k/p) \leq 13(3 + 3k/d)$$

generalized elementary matrices.

Combining lemmas 1.2 and 1.3 we obtain:

Theorem 1.4 *Let H be a normal subgroup of finite index in $G_{d,k}$. Then every matrix in $G_{d,k} = \text{SL}_d(\mathbb{Z}[x_1, \dots, x_k])$ can be written as a product of $77 + 39k/d$ generalized elementary matrices modulo H .*

Finally we can complete the proof of Theorem 6. Let $g \in G_{d,k}$, using theorem 1.4 we can write g as a product of $77 + 39k/d$ GEMs v_s and an element in $h \in H = \ker \rho$. As in the proof of Theorem 5 this implies that

$$\|\rho(g)v - v\| \leq \|\rho(h)v - v\| + \sum_{s=1}^{77+39k/d} \|\rho(v_s)v - v\| < 2(77 + 39k/d)M_d(k)\epsilon.$$

If $T_{d,k} = [\sqrt{2}(77 + 39k/d)M_d(k)]^{-1}$ then we will have that $\|\rho(g)v - v\| < \sqrt{2}$ for any $g \in G$ and the representation ρ will have an invariant vector, which completes the proof that the group $\mathrm{SL}_d(R)$ has the property T and also provides a lower bound for the τ -constant:

$$\tau(G_{d,k}, S_{d,k}) \geq \frac{1}{\sqrt{2}(77 + 39k/d)M_d(k)}.$$

It is easy to see that we have the inequality

$$\sqrt{2}(77 + 39k/d)M_d(k) < 800\sqrt{d} \left(1 + (k/d)^{3/2}\right),$$

which completes the proof of theorem 6. $\square$

Theorems 5 and 1.4 are now proved Lemmas 1.1 and 1.2.

2 Relative property T of $\mathrm{EL}_2(R) \ltimes R^2, R^2$

In this section we shall show that the pair of groups $(\mathrm{EL}_2(R) \ltimes R^2, R^2)$ has relative property T , and we shall estimate its relative Kazhdan constant. This result was first proven by M. Burger [2] in the case $R = \mathbb{Z}$ and later generalized by Y. Shalom [14] to the case of finitely generated commutative rings. In their proofs they use that $\mathrm{SL}_2(\mathbb{Z})$ is generated by the elementary matrices (and is thus equal to $\mathrm{EL}_2(\mathbb{Z})$), but this assumption is not necessary. The argument which follows gives a better estimate of the relative Kazhdan constant in the case of commutative ring R and is based on ideas from [7].

First we start with a definition of the *relative property T* for finitely generated discrete groups:

Definition 2.1 *A pair of groups (G, H) such that $H \leq G$, where G is generated by a finite set S is said to have the Relative Kazhdan Property T if there is $\epsilon > 0$ such that for every irreducible (continuous) unitary representation $\rho : G \rightarrow U(\mathcal{H})$ on a Hilbert space $\mathcal{H}$ without H invariant vectors and for every vector $v \neq 0$ in $\mathcal{H}$ there is some $s \in S$ such that $\|\rho(s)v - v\| > \epsilon\|v\|$. The largest ϵ with this property is called the Relative Kazhdan constant and is denoted by $\mathcal{K}(G, H; S)$. It is easy to see that the relative property T depends only on the groups G and H and is independent of the generating set S , however the value of the relative Kazhdan constant depends also on S .*

We will show that $(\mathrm{EL}_2(R_k^n) \ltimes R_k^{n2}, R_k^{n2})$ has the relative property T by proving the following theorem (which is inspired by [14]):

Theorem 2.2 *Let F_1 be set of $4(k+1)$ elementary matrices in $\mathrm{EL}(2; R_k^n)$ with ± 1 and $\pm x_j$ off the diagonal and F_2 be the set of standard basis vectors in R_k^{n2} and their inverses. Let $\rho : \Gamma \rightarrow U(\mathcal{H})$ be a unitary representation of the group $\Gamma := \mathrm{EL}_2(R_k^n) \ltimes R_k^{n2}$ in a Hilbert space $\mathcal{H}$. Let $v \in \mathcal{H}$ be a unit vector such that*

$\|\rho(g)v - v\| < \epsilon = M(k)^{-1}$ for all $g \in F_1 \cup F_2$, then $\mathcal{H}$ contains an $R_k^{n^2}$ -invariant vector. Here the numbers $M(k)$ are defined by $M(0) = 2 + \sqrt{10}$ and

$$M(k)^2 = 4M(0)^2 + (6\sqrt{k} + 4)\sqrt{(M(0) + \sqrt{k})^2 + k + 2\sqrt{k} + 1} + 6\sqrt{k}M(0) + 7k + 10\sqrt{k} + 2.$$

This shows that $\text{EL}_2(R_k^n) \ltimes R_k^{n^2}, R_k^{n^2}$ has relative property T and

$$\mathcal{K}(\text{EL}_2(R_k) \ltimes R_k^2, R_k^2; F_1 \cup F_2) \leq 1/M(k).$$

Using the definition of $M(k)$ it can be verified that $M(0) < 5.17$, $M(1) < 14.92$, $M(2) < 16.47$ and $M(k) \leq \sqrt{18}(\sqrt{k} + 3)$.

Before starting the proof of this theorem, let us note that there are two natural actions of $\text{EL}_2(R)$ on R^2 , a ‘left’ action where we think of the elements in R^2 as column vectors, i.e., we embed $\text{EL}_2(R) \ltimes R^2$ in $\text{EL}_3(R)$ as follows

$$(A, B) \rightarrow \begin{pmatrix} A & B \\ 0 & 1 \end{pmatrix}.$$

There is also a ‘right’ one where we think of R^2 as a row vectors and we use the embedding

$$(A, B) \rightarrow \begin{pmatrix} A & 0 \\ B & 1 \end{pmatrix}.$$

In the case of commutative ring these two actions give isomorphic semidirect products, which is not always the case if the ring R is not commutative. However we have that the ‘left’ semi-direct product is isomorphic to the ‘right’ semidirect product over the opposite ring R^{opp} . In the proof we will consider only the ‘left’ action. By the previous remark the theorem is valid also for the ‘right’ semidirect products and the proof is essentially the same.

Proof of Theorem 2.2: Assume that we have a representation ρ which satisfies all the conditions in the theorem and the Hilbert space $\mathcal{H}$ does not contain an R^2 invariant vector. Let P be the projection valued measure on the dual $(R^2)^* = R^{*2}$, coming from the restriction of the representation ρ to R^2 . Here by R^* we have denoted the dual of the additive group of the ring R , i.e., $R^* = \text{Hom}(R, \mathbb{C})$, where $S^1 = \{z \in \mathbb{C} \mid |z| = 1\} = U(\mathbb{C})$ is the unit circle in the complex plane. The set R^* is an abelian group with natural topology. For $t \in R^*$ and $r \in R$ by $t(r)$ we will denote the natural pairing of R^* and R . Let μ_v be the probability measure on R^{*2} , defined by $\mu_v(B) = \langle P(B)v, v \rangle$. The measure μ_v is supported on $R^{*2} \setminus \{(0, 0)\}$, because by assumption $\mathcal{H}$ does not contain an R^2 invariant vector and by the construction of the measure P , $P(\{0\})$ is the projection onto the subspace of R^2 invariant vectors in $\mathcal{H}$.

For an element $T \in R^{*2}$ we will write $T = (t_1, t_2)^t$, where t_1 and t_2 are in R^* .

Lemma 2.3 *Let $P_i = \{T \in R^{*2} \mid \Re(t_i(1)) < 0\}$, then $\mu_v(P_i) \geq 1 - \epsilon^2/2$.*

Proof: By the definition of the measure μ_v , we have

$$\|\rho(g_i)v - v\|^2 = \int_{R^{*2}} |t_i(1) - 1|^2 d\mu_v,$$

where $g_1, g_2 \in G$ form the standard basis of R^2 . By assumption v is almost preserved by $\rho(g_i)$ therefore $\|\rho(g_i)v - v\|^2 \leq \epsilon$. If we break the integral into two integrals over P_i and its complement, we get

$$\epsilon^2 \geq \int_{R^{*2} \setminus P_i} |t_i(1) - 1|^2 d\mu_v + \int_{P_i} |t_i(1) - 1|^2 d\mu_v \geq \int_{R^{*2} \setminus P_i} 2 d\mu_v = 2(1 - \mu_v(P_i)),$$

which gives that the measure of the set P_i is large enough. $\square$

Lemma 2.4 *For every measurable set $B \subset R^{*2}$ and every elementary matrix $g \in F_1$, we have that*

$$|\mu_v(gB) - \mu_v(B)| \leq 2\epsilon\sqrt{\mu_v(B)} + \epsilon^2 \quad \text{or} \quad |\sqrt{\mu_v(gB)} - \sqrt{\mu_v(B)}| \leq \epsilon.$$

*The action of $\text{EL}_2(R)$ on R^{*2} is the dual of the action of $\text{EL}_2(R)$ on R^2 .*

Proof: Using the properties of the projection valued measure P , we have

$$\begin{aligned} |\mu_v(gB) - \mu_v(B)| &= |\langle \pi(g^{-1})P(B)\pi(g)v, v \rangle - \langle P(B)v, v \rangle| \leq \\ &\leq |\langle \pi(g^{-1})P(B)(\pi(g)v - v), v \rangle| + |\langle P(B)v, (\pi(g)v - v) \rangle| = \\ &= 2|\langle \pi(g)v - v, P(B)v \rangle| + \langle P(B)(\pi(g)v - v), \pi(g)v - v \rangle \leq \\ &\leq 2\epsilon\sqrt{\mu_v(B)} + \epsilon^2, \end{aligned}$$

where the final inequality follows from the facts that v is (F, ϵ) invariant vector and $\|P(B)v\|^2 = \mu_v(B)$. $\square$

We will need the following slight generalization of the above lemma:

Lemma 2.5 *Let A and B be measurable sets in R^{*2} . Suppose that A decomposes as a disjoint union of the sets A_i for $i = 1, \dots, s$ and there exist elements $g_i \in F_1$ such that the sets $B_i = g_i(A_i)$ are disjoint subsets of B then*

$$\sqrt{\mu_v(A)} \leq \sqrt{\mu_v(B)} + \sqrt{s}\epsilon.$$

Proof: Applying lemma 2.4 to the sets A_i yields to

$$\begin{aligned} \mu_v(A) &= \sum_{i=1}^s \mu_v(A_i) \leq \sum_{i=1}^s \left[\mu_v(B_i) + 2\epsilon\sqrt{\mu_v(B_i)} + \epsilon^2 \right] \leq \\ &\leq \sum_{i=1}^s \mu_v(B_i) + 2\epsilon\sqrt{s \sum_{i=1}^s \mu_v(B_i)} + s\epsilon^2 \leq \\ &\leq \mu_v(B) + 2\epsilon\sqrt{s\mu_v(B)} + s\epsilon^2 = \left(\sqrt{\mu_v(B)} + \sqrt{s}\epsilon \right)^2, \end{aligned}$$

where we have used that for nonnegative numbers a_i we have the inequality

$$\sqrt{\sum_{i=1}^s a_i} \leq \sum_{i=1}^s \sqrt{a_i} \leq \sqrt{s \sum_{i=1}^s a_i}. \square$$

Let us first consider the case when $k = 0$ and $R = \mathbb{Z}$. In this case the dual of the ring R is just the unit circle in complex plane. Also we can think of R^* as the torus $\mathbb{T}^2 = S^1 \times S^1 = \mathbb{R}^2/\mathbb{Z}^2$ which we will identify with the square $(-1/2, 1/2] \times (-1/2, 1/2]$. It is well known that there are only a few $\text{SL}_2(\mathbb{Z})$ invariant measures on the torus $\mathbb{T}^2$. The next lemma is a quantitative version of this fact.


Lemma 2.6 *Let μ be a finitely additive measure on $\mathbb{T}^2$ such that:*

1. $\mu(|x| \geq 1/4) \leq \epsilon^2/2$ and $\mu(|y| \geq 1/4) \leq \epsilon^2/2$,
2. $|\mu(gB) - \mu(B)| \leq 2\epsilon\sqrt{\mu(B)} + \epsilon^2$ for any Borel set B and any elementary matrix $g \in F_1 \cap \text{SL}_2(\mathbb{Z})$.

Then the measure μ satisfies:

$$\mu(\mathbb{T}^2 \setminus \{(0, 0)\}) \leq (2 + \sqrt{10})^2 \epsilon^2.$$

Proof: Let us define the Borel subsets A_i and A'_i of $\mathbb{T}^2$ using the picture:



Each set A_i or A'_i consists of the interiors of two triangles and part of their boundary (not including the vertices). The sets A_i do not contain the side which is part of the small square, they also do not contain their clockwise boundary but contain the counter-clockwise one. Each set A'_i includes only the part of its boundary which lies on the small square.

It can be seen, from the picture, that the elementary matrices $e_{ij}^\pm = e_{ij}(\pm 1) \in F_1$, act on the sets A_i as follows:

$$\begin{aligned} e_{12}^+(A_3 \cup A'_4) &= A_3 \cup A_4 & e_{21}^+(A'_3 \cup A_4) &= A_3 \cup A_4 \\ e_{12}^-(A'_1 \cup A_2) &= A_1 \cup A_2 & e_{21}^-(A_1 \cup A'_2) &= A_1 \cup A_2. \end{aligned}$$

Using the properties of the measure μ the above equalities imply the inequalities:

$$\begin{aligned}\mu(A_1) + \mu(A_2) &\leq \mu(A'_1) + \mu(A_2) + \epsilon^2 + 2\epsilon\sqrt{\mu(A'_1) + \mu(A_2)} \\ \mu(A_1) + \mu(A_2) &\leq \mu(A_1) + \mu(A'_2) + \epsilon^2 + 2\epsilon\sqrt{\mu(A_1) + \mu(A'_2)} \\ \mu(A_3) + \mu(A_4) &\leq \mu(A'_3) + \mu(A_4) + \epsilon^2 + 2\epsilon\sqrt{\mu(A'_3) + \mu(A_4)} \\ \mu(A_3) + \mu(A_4) &\leq \mu(A_3) + \mu(A'_4) + \epsilon^2 + 2\epsilon\sqrt{\mu(A_3) + \mu(A'_4)}.\end{aligned}$$

Adding these inequalities and noticing that

$$\begin{aligned}\mu(A'_1) + \mu(A'_4) &\leq \mu(\{|x| \geq 1/4\}) \leq \epsilon^2/2 \quad \text{and} \\ \mu(A'_2) + \mu(A'_3) &\leq \mu(\{|y| \geq 1/4\}) \leq \epsilon^2/2\end{aligned}$$

we obtain

$$\begin{aligned}\sum \mu(A_i) &\leq 4\epsilon^2 + \sum \mu(A'_i) + 2\epsilon\sqrt{4\left(\sum \mu(A_i) + \sum \mu(A'_i)\right)} \leq \\ &\leq 5\epsilon^2 + 4\epsilon\sqrt{\sum \mu(A_i) + \epsilon^2}.\end{aligned}$$

After substituting $c = \sqrt{\sum \mu(A_i) + \epsilon^2}$ and solving the resulting quadratic inequality one obtains $\sum \mu(A_i) \leq (13 + 4\sqrt{10})\epsilon^2$. Finally we can use that

$$\begin{aligned}\mu(\mathbb{T}^2 \setminus \{(0,0)\}) &\leq \sum \mu(A_i) + \mu(\{|x| \geq 1/4\}) + \mu(\{|y| \geq 1/4\}) \leq \\ &\leq (14 + 4\sqrt{10})\epsilon^2 = (2 + \sqrt{10})^2\epsilon^2,\end{aligned}$$

which completes the proof of the lemma. $\square$

This completes the proof of Theorem 2.2 in the case $k = 0$ — by assumption the Hilbert space $\mathcal{H}$ does not contains R^2 invariant vectors. However, Lemma 2.6 gives us that $\mu(0,0) > 0$ if $\epsilon < \frac{1}{2+\sqrt{10}} = M(0)^{-1}$ which is a contradiction with the assumption the $\mathcal{H}$ does not contain $\mathbb{Z}^2$ invariant vectors.

In order to prove the general case of Theorem 2.2 we need to generalize Lemma 2.6 to the case of larger rings. Recall, that R_k^n denotes the free ring $\mathbb{Z}\langle x_1, \dots, x_k \rangle$. Let F denote the set of elementary matrices in $\text{EL}_2(R_k^n)$, with ± 1 and $\pm x_i$ off the diagonal. Denote the dual of the abelian group $R_k^n = \mathbb{Z}\langle x_1, \dots, x_k \rangle$ by R_k^{n*} .

Lemma 2.7 *Let μ be a finitely additive measure on R_k^{n*2} , such that*

1. $\mu(\{T \mid \Re(t_1(1)) < 0\}) \leq \epsilon^2/2$ and $\mu(\{T \mid \Re(t_2(1)) < 0\}) \leq \epsilon^2/2$. Here t_i denote the components of the element $T \in R_k^{n*2}$.
2. $|\mu(gB) - \mu(B)| \leq 2\epsilon\sqrt{\mu(B)} + \epsilon^2$ for any Borel set $B \subset R_k^{n*2}$ and any elementary matrix $g \in F_1 \subset \text{EL}_2(R_k)$.

Then we have

$$\mu(R_k^{n*2} \setminus \{(0,0)\}) \leq M(k)^2\epsilon^2,$$

where the number $M(k)$ is defined in Theorem 2.2.

Proof: The idea for the proof is based on the proof of Lemma 3.3 from [14] - the main difference is that we do not use induction on k and do everything in one step, which allows us to drop the assumption that the ring R is commutative.

By the Lemma 2.6 we know that if we only consider the action of $\mathbb{Z} \subset R_k^n$ we have that

$$\mu(\mathbb{Z}^{*2} \setminus \{(0, 0)\}) \leq M(0)^2 \epsilon^2.$$

There is a natural restriction map $\text{res} : R_k^{n*} \rightarrow \mathbb{Z}^*$ coming from the inclusion $\mathbb{Z} \subset R_k^n$ and the correct way of writing the above inequality is

$$\mu(R_k^{n*2} \setminus \text{res}^{-1}\{(0, 0)\}) \leq M(0)^2 \epsilon^2.$$

By definition, we have that $R_k^n = \mathbb{Z}\langle x_1, \dots, x_k \rangle$, which allows us to write

$$R_k^{n*} \simeq \mathbb{Z}^*\langle x_1^{-1}, \dots, x_k^{-1} \rangle$$

as a topological space. Here $\mathbb{Z}^*\langle x_1^{-1}, \dots, x_k^{-1} \rangle$ is the formal power series on non commuting variables x_k^{-1} with coefficients in $\mathbb{Z}^*$. This isomorphism gives a valuation ν on R_k^{n*} , corresponding to the total degree of the leading term.

Let us define the following subsets of $R_k^{n*2} \setminus \{0, 0\}$:

$$\begin{aligned} A &= \{(\chi_1, \chi_2) \mid \nu(\chi_1) > \nu(\chi_2) > 0\} \\ B &= \{(\chi_1, \chi_2) \mid \nu(\chi_1) = \nu(\chi_2) > 0\} \\ C &= \{(\chi_1, \chi_2) \mid \nu(\chi_2) > \nu(\chi_1) > 0\} \\ D &= \{(\chi_1, \chi_2) \mid \nu(\chi_1) = 0 \text{ or } \nu(\chi_2) = 0\} \end{aligned}$$

Lets consider the action of $e_{i,j}(r), r \in R_k^n$ on the element $(\chi_1, \chi_2) \in R_k^{n*2}$, by the definition of this action we have

$$(e_{1,2}(r) \circ (\chi_1, \chi_2)) \begin{pmatrix} r_1 \\ r_2 \end{pmatrix} = (\chi_1, \chi_2) \left(e_{1,2}(r)^{-1} \begin{pmatrix} r_1 \\ r_2 \end{pmatrix} \right) = (\chi_1, \chi_2) \begin{pmatrix} r_1 - r \cdot r_2 \\ r_2 \end{pmatrix}$$

and

$$(e_{2,1}(r) \circ (\chi_1, \chi_2)) \begin{pmatrix} r_1 \\ r_2 \end{pmatrix} = (\chi_1, \chi_2) \left(e_{2,1}(r)^{-1} \begin{pmatrix} r_1 \\ r_2 \end{pmatrix} \right) = (\chi_1, \chi_2) \begin{pmatrix} r_1 \\ r_2 - r \cdot r_1 \end{pmatrix}$$

This shows that

$$e_{1,2}(1)(A) \subset B \quad e_{2,1}(1)(C) \subset B.$$

and the first two inequalities can be written as

$$\sqrt{\mu(A)} \leq \sqrt{\mu(B)} + \epsilon \quad \text{and} \quad \sqrt{\mu(C)} \leq \sqrt{\mu(B)} + \epsilon.$$

Claim: The union $A \cup B$ can be decomposed as a disjoint union of k sets A_i such that the sets $\widetilde{A}_i = e_{1,2}(x_i)(A_i)$ are disjoint and lie in $C \cup D$. Similarly, $C \cup B$ can be written as $\cup C_i$ such that $\widetilde{C}_i = e_{2,1}(x_i)(C_i)$ are disjoint subsets of $A \cup D$.

Proof of the Claim: Let $(\chi_1, \chi_2) \in A \cup B$. Let $\bar{\chi}$ be the leading term of χ_2 . Then $\bar{\chi}$ is a non trivial element of the dual of the space of all homogeneous

polynomials of degree $\nu(\chi_2) > 0$ in $x_1, \dots, x_k$. Let $\bar{\chi}_{(i)}$ be the element in the dual of the space of all homogeneous polynomials of degree $\nu(\chi_2) - 1$ defined by

$$\bar{\chi}_{(i)}(f) = \bar{\chi}(x_i f).$$

Let $s(\bar{\chi})$ be the smallest index such that $\bar{\chi}_{(i)}$ is not trivial – such index exists because $\bar{\chi}$ is non zero. Define the sets A_i by

$$A_i = \{(\chi_1, \chi_2) \in A \cup B \mid s(\bar{\chi}) = i\}.$$

It is clear that $A \cup B$ is a disjoint union of A_i and that $\widetilde{A_i} = e_{1,2}(x_i)(A_i)$ are disjoint subsets of $C \cup D$.

The second part of the claim is proved the same way but one uses the first component χ_1 of $\bar{\chi}$ not the second one χ_2 . $\square$

We are ready to complete the proof of Lemma 2.7. Applying Lemma 2.5 for the sets $A \cup B$ and $C \cup D$ gives us:

$$\begin{aligned} \sqrt{\mu(A \cup B)} &\leq \sqrt{\mu(C \cup D)} + \sqrt{k}\epsilon \\ \sqrt{\mu(C \cup D)} &\leq \sqrt{\mu(A \cup B)} + \sqrt{k}\epsilon. \end{aligned}$$

Squaring and adding the the above inequalities leads to

$$\begin{aligned} 2\mu(B) &\leq 2\mu(D) + 2\epsilon\sqrt{k(\mu(A) + \mu(D))} + 2\epsilon\sqrt{k(\mu(A) + \mu(D))} + 2k\epsilon^2 \leq \\ &\leq 2\mu(D) + 4\epsilon\sqrt{k\mu(D)} + 2\epsilon\sqrt{k\mu(A)} + 2\epsilon\sqrt{k\mu(C)} + 2k\epsilon^2 \leq \\ &\leq 2\mu(D) + 4\epsilon\sqrt{k\mu(D)} + 4\epsilon\sqrt{k\mu(B)} + (2k + 4\sqrt{k})\epsilon^2 \end{aligned}$$

The last inequality can be rewritten as

$$(\sqrt{\mu(B)} - \sqrt{k}\epsilon)^2 \leq (\sqrt{\mu(D)} + \sqrt{k}\epsilon)^2 + (k + 2\sqrt{k})\epsilon^2$$

Therefore we have

$$\begin{aligned} \sqrt{\mu(A)} &\leq \left(\sqrt{(M(0) + \sqrt{k})^2 + (k + 2\sqrt{k})} + \sqrt{k} + 1 \right) \epsilon \\ \sqrt{\mu(B)} &\leq \left(\sqrt{(M(0) + \sqrt{k})^2 + (k + 2\sqrt{k})} + \sqrt{k} \right) \epsilon \\ \sqrt{\mu(C)} &\leq \left(\sqrt{(M(0) + \sqrt{k})^2 + (k + 2\sqrt{k})} + \sqrt{k} + 1 \right) \epsilon \end{aligned}$$

Finally we have

$$\begin{aligned} \mu(R_k^{*2} \setminus \{(0, 0)\}) &= \mu(A \cup B \cup C \cup D) \leq \mu(A) + \mu(B) + \mu(C) + \mu(D) \leq \\ &\leq \left(4M(0)^2 + (6\sqrt{k} + 4)\sqrt{(M(0) + \sqrt{k})^2 + k + 2\sqrt{k}} + \right. \\ &\quad \left. 6\sqrt{k}M(0) + 7k + 10\sqrt{k} + 2 \right) \epsilon^2 = M(k)^2 \epsilon^2, \end{aligned}$$

which completes the proof of lemma 2.7. $\square$

Finally Theorem 2.2 follows easily: by assumption the Hilbert space $\mathcal{H}$ does not contain R^2 invariant vectors. However lemma 2.7 gives us that $\mu(0, 0) > 0$ if $\epsilon < M(k)^{-1}$ which is a contradiction.

Using that $M(0) = 2 + \sqrt{10}$, it can be easily shown that

$$M(0) < 5.17, \quad M(1) < 14.92, \quad M(2) < 16.47, \quad \text{and} \quad M(k) < 3\sqrt{2}(\sqrt{k}+3). \square$$

Corollary 2.8 *Let $\rho : \Gamma \rightarrow U(\mathcal{H})$ be a unitary representation of the group*

$$\Gamma = \text{EL}_2(R_k^n) \ltimes R_k^{n^2}.$$

Let $v \in \mathcal{H}$ be a unit vector such that $\|\rho(g)v - v\| < \epsilon$ for $g \in F_1 \cup F_2$. Then for every g in $R_k^{n^2}$ we have $\|\rho(g)v - v\| \leq 2M(k)\epsilon$.

Proof: Let us decompose the Hilbert space $\mathcal{H}$ as the orthogonal direct sum $\mathcal{H}_0 \oplus \mathcal{H}_1$, where $\mathcal{H}_0$ contains all $R_k^{n^2}$ invariant vectors and $\mathcal{H}_1 = \mathcal{H}_0^\perp$. We have that both $\mathcal{H}_0$ and $\mathcal{H}_1$ are closed under the action of the group Γ , because $R_k^{n^2}$ is a normal subgroup of Γ .

Let us write $v = v_0 + v_1$, where $v_i \in \mathcal{H}_i$. Since there are no $R_k^{n^2}$ -invariant vectors in $\mathcal{H}_1$, there exists $h \in F \cup G$ such that $\|\rho(h)v_1 - v_1\| \geq M(k)^{-1}\|v_1\|$. But, we have that

$$\|\rho(h)v - v\|^2 = \|\rho(h)v_0 - v_0\|^2 + \|\rho(h)v_1 - v_1\|^2 \leq \epsilon^2.$$

The final inequality implies $\|v_1\| \leq M(k)\epsilon$. For any $g \in R_k^{n^2}$, we have

$$\|\rho(g)v - v\|^2 = \|\rho(g)v_0 - v_0\|^2 + \|\rho(g)v_1 - v_1\|^2 \leq 0 + 4\|v_1\|^2 \leq 4M(k)^2\epsilon^2,$$

therefore $\|\rho(g)v - v\| \leq 2M(k)\epsilon$. $\square$

As another corollary we obtain part a) of Lemma 1.1 from Section 1.

Using the similar methods we can estimate the relative Kazhdan constant for the pairs

$$\text{EL}_p(R) \ltimes R^p, R^p,$$

with $p \geq 2$ and

$$(\text{EL}_p(R) \times \text{EL}_q(R)) \ltimes R^{pq}, R^{pq},$$

where $p \geq 2$. Here we consider the elements of $(\text{EL}_p(R) \times \text{EL}_q(R)) \ltimes R^{pq}$ as $(p+q) \times (p+q)$ matrices over R of type

$$\begin{pmatrix} A & C \\ 0 & B \end{pmatrix},$$

where $A \in \text{EL}_p(R)$, $B \in \text{EL}_q(R)$ and $C \in R^{pq}$. Equivalent way of say this is the we consider R^{pq} as a left $\text{EL}_p(R)$ -module and a right $\text{EL}_q(R)$ -module.

Theorem 2.9 a) Let $F_{p;k}$ be the following generating set of $\Gamma_p = \text{EL}_p(R_k^n) \ltimes R_k^{n,p}$

$$F_{p;k} = \{\text{Id} \pm e_{ij} \in \text{EL}_p(R) \mid 1 \leq i \neq j \leq p\} \cup \\ \cup \{\text{Id} \pm x_l \cdot e_{ij} \in \text{EL}_p(R) \mid |i - j| = 1, 1 \leq l \leq k\} \cup \\ \cup \{\pm 1_i \in R^p \mid 1 \leq i \leq p\}$$

consisting of some elementary matrices and the standard generators of R^p .

Let $\rho : \Gamma_p \rightarrow U(\mathcal{H})$ be a unitary representation of Γ_p acting on a Hilbert space $\mathcal{H}$. Let $v \in \mathcal{H}$ be a unit vector such that $\|\rho(g)v - v\| < \epsilon = M(k;p)^{-1}$ for all $g \in F_p$, then $\mathcal{H}$ contains an R_k^p -invariant vector. Here $M(k;p)$ denotes the number

$$M(k;p) = \sqrt{2M(k)^2 + 2M(k)\sqrt{p-2} + p-2},$$

where $M(k)$ is the constant defined in Theorem 2.2.

b) Let $F_{p,q;k}$ be the generating set of $\Gamma_{p,q} = (\text{EL}_p(R_k^n) \times \text{EL}_q(R_k^n)) \ltimes R_k^{n,pq}$: consisting of the elementary matrices in $\text{EL}_p(R)$ and $\text{EL}_q(R)$ and the standard generators of R^{pq} .

Let $\rho : \Gamma_{p,q} \rightarrow U(\mathcal{H})$ be a unitary representation of $\Gamma_{p,q}$ on a Hilbert space $\mathcal{H}$. Let $v \in \mathcal{H}$ be a unit vector such that $\|\rho(g)v - v\| < \epsilon = M(k;p,q)^{-1}$ for all $g \in F_{p,q}$, then $\mathcal{H}$ contains an H -invariant vector. Here $M(k;p,q)$ denotes the number

$$M(k;p,q) = \sqrt{2M(k;p)^2 + 2M(k;p)\sqrt{q-1} + q-1}.$$

Proof: a) The proof is similar to the one of Theorem 2.2. Let μ_v be the measure on R_k^{*p} coming from the restriction of the representation ρ to the abelian group R_k^p . As in the case $p = 2$ the measure μ_v is almost preserved under the action of $\text{EL}_p(R)$.

We will write an element $T \in R_k^{*p}$ as $(t_1, \dots, t_p)^t$, where $t_i \in R_k^*$. Using the restriction $R_k^{*p} \rightarrow R_k^{*2}$ coming from the inclusion $R_k^2 \subset R_k^p$ and Lemma 2.8 we can see that

$$\mu_v(\{T \mid t_1 \neq 0 \text{ or } t_2 \neq 0\}) \leq M(k)\epsilon^2.$$

Let us define the Borel subsets B_i, C_i of R_k^{*p} by

$$B_i = \{T \mid t_k = 0 \text{ for } k \leq i\}, \text{ and} \\ C_i = \{T \mid t_1 = t_i \neq 0, t_k = 0 \text{ for } 1 < k < i\}.$$

The elementary matrix $g_{1i} \in \text{SL}_p \subset \text{EL}_p(R)$ sends $B_{i-1} \setminus B_i$ into C_i for any $i \geq 3$. Now, notice that the sets C_i are disjoint for $i = 2, \dots, p$, and their union lies in the set

$$C := \{T \mid t_1 \neq 0, t_2 = 0\} \subset \{T \mid t_1 \neq 0 \text{ or } t_2 \neq 0\}.$$

Therefore, by lemma 2.5 we have

$$\sqrt{\mu_v(B_2 \setminus B_p)} \leq \sqrt{\mu_v(C)} + \sqrt{(p-2)\epsilon}.$$

We know that the measure of C is very small which gives

$$\begin{aligned}\mu_v(R_k^{*p} \setminus \{(0, \dots, 0)\}) &= \mu_v(R_k^{*p} \setminus B_2) + \mu_v(B_2 \setminus B_p) \leq \\ &\leq M(k)^2 \epsilon^2 + \left(M(k)\epsilon + \sqrt{p-2}\epsilon\right)^2 = M(k;p)^2 \epsilon^2.\end{aligned}$$

By assumption $\epsilon < M(k;p)^{-1}$ which implies that the measure of the zero in R_k^{*p} is $\mu_v(\{(0, \dots, 0)\}) > 0$, and this shows that $\mathcal{H}$ contains R^p -invariant vectors.

b) The idea is the same as the one in part a): Let μ_v be the measure on R_k^{*pq} coming from the restriction of the representation ρ to the abelian group R_k^{pq} . As in the case $q = 1$ the measure μ_v is almost invariant under the action of $\text{EL}_p(R)$ and $\text{EL}_q(R)$.

We will write an element $T \in R_k^{*pq}$ as $(T_1, \dots, T_q)$ where $T_i \in R_k^{*p}$. Using the restriction $R_k^{*pq} \rightarrow R_k^{*p}$ coming from the inclusion $R_k^p \subset R_k^{pq}$ which is given by $T \rightarrow T_1$, and part a) we have

$$\mu_v(\{T \mid T_1 \neq 0\}) \leq M(k;p)^2 \epsilon^2.$$

Define the Borel subsets B_i, C_i of R_k^{*pq} by

$$\begin{aligned}B_i &= \{T \mid T_k = 0 \text{ for } k \leq i\}, \text{ and} \\ C_i &= \{T \mid T_1 = T_i \neq 0, T_k = 0 \text{ for } 1 < k < i\}.\end{aligned}$$

The elementary matrix $g_{1i} \in \text{EL}_q$ sends $B_{i-1} \setminus B_i$ into C_i for any i . Therefore, we have

$$\mu_v(B_{i-1} \setminus B_i) \leq \mu_v(C_i) + \epsilon^2 + 2\epsilon\sqrt{\mu_v(C_i)}.$$

The sets C_i , for $i = 1, \dots, q$, are disjoint and their union lies in the set

$$C = \{T \mid T_1 \neq 0\}.$$

Therefore, by lemma 2.5 we have

$$\sqrt{\mu_v(B_1 \setminus B_q)} \leq \sqrt{\mu_v(C)} + \sqrt{q-1}\epsilon.$$

We know that the measure of C is very small which gives us

$$\begin{aligned}\mu_v(R_k^{*pq} \setminus \{0\}) &= \mu_v(R_k^{*pq} \setminus B_1) + \mu_v(B_1 \setminus B_q) \leq \\ &\leq M(k;p)^2 \epsilon^2 + \left(M(k;p)\epsilon + \sqrt{q-1}\epsilon\right)^2 = M(k;p,q)^2 \epsilon^2\end{aligned}$$

By assumption $\epsilon < M(k;p,q)^{-1}$ which implies that $\mu_v(\{0\}) > 0$. This shows that $\mathcal{H}$ contains R^{pq} -invariant vectors. $\square$

Corollary 2.10 *Let $\rho : G \rightarrow U(\mathcal{H})$ be a unitary representation of the group $G = \text{EL}(d; R)$. Let $v \in \mathcal{H}$ be a unit vector such that $\|\rho(g)v - v\| < \epsilon$ for $g \in F_1 \cup F_2$, where F_1 is the set of $2(d^2 - d)$ elementary matrices with ± 1 off the diagonal and F_2 is the set of $4k(d-1)$ elementary matrices $\text{Id} \pm x_s e_{ij}$ with $|i-j| = 1$. Then for every generalized elementary matrix g we have $\|\rho(g)v - v\| \leq 2M_d(k)\epsilon$, where $M_d(k) = 2M(k) + \sqrt{3d} \leq 6\sqrt{2}(\sqrt{k} + 3) + \sqrt{3d}$.*

Proof: Use the inequality

$$M(k; p, q) \leq 2M(k) + \sqrt{2p} + \sqrt{q} \leq 2M(k) + \sqrt{3d} = M_d(k) \text{ if } p + q \leq d. \square$$

This corollary immediately implies part b) of Lemma 1.1 from Section 1.

3 Uniform bounded generation of $\mathrm{SL}_d(\bar{R})$

Lemma 1.2 is a well-known result for fields. In the case of finite commutative rings the same proof works. Almost everything works also in the case of ring satisfying Bass stable range condition.

Definition 3.1 *A ring R is said to satisfy the Bass stable range condition if for any elements $a, b \in R$ such that $Ra + Rb = R$, there exist $t \in R$ such that $R(a + tb) = R$.*

There is an equivalent definition: *If a be an element R and I be a left ideal R , such that the left ideal generated by a and I coincides with R . Then there exist $i \in I$ such that the left ideal generated by $u = a + i$ is the whole ring R .*

This is the left Bass stable range condition, there is also a right version and it is known that the two conditions are equivalent. Any finite ring satisfies the Bass stable range condition see [17].

If a ring R satisfies the Bass stable range condition, there is an algorithm which writes an element in $\mathrm{EL}_d(R)$ as a product of small number of elementary matrices and an element in $\mathrm{EL}_{d-1}(R)$.

Lemma 3.2 *Let the ring $\bar{R}$ satisfied the Bass stable range condition. Then any element $g \in \mathrm{EL}_d(\bar{R})$ can be written as*

$$g = \prod_{s=1}^{d^2-1} E_s \cdot g_{11} \cdot \prod_{t=1}^{(d^2-d)/2} E'_t$$

where each $E_s, E'_t \in E(\bar{R})$ is an elementary matrix and g_{11} lies in the copy of $\mathrm{GL}_1(\bar{R})$ embedded in the upper right corner, i.e., $g_{11} = e_{11}(a)$, where a is a unit in $\bar{R}$.

Proof: Since the ring $\bar{R}$ satisfies the Bass stable range condition Lemma 3.2 is a consequence of the familiar argument that a matrix $g \in \mathrm{SL}_d(K)$ can be reduced by successive applications of row and column operation to the identity matrix: Each of these operations is in fact a multiplication by an elementary matrix from left or right.

The following well-known algorithm produces such decomposition (cf. [5], Remark 10):

Let $g \in \mathrm{GL}_d(\bar{R})$. Let $\bar{I}$ be the left ideal in $\bar{R}$ generated by $g_{2d}, \dots, g_{dd}$ in the last column of g . The matrix g is left invertible therefore the left ideal generated

by g_{1d} and $\bar{I}$ coincide with the ring $\bar{R}$. The Bass stable range condition implies that there exist $i \in \bar{I}$ such that $g_{1d} + i$ is a unit in $\bar{R}$. This allows us to make last entry on the first row is an invertible element in $\bar{R}$ by multiplying with $d - 1$ elementary matrices to the left. Using an extra multiplication to the left we can make the d, d entry equal to 1, next with $d - 1$ left and $d - 1$ right multiplications by elementary matrices we can transform the matrix g to an element in $\text{GL}_{d-1} \times 1$ (sitting in the top left corner). Thus the reduction from GL_d to GL_{d-1} can be done using $3d - 2$ elementary matrices, and by induction using $(3d^2 - d - 2)/2$ we can transform any matrix to an element in GL_1 .

If we begin with an element in EL_d then it is clear that after applying the above algorithm the resulting element lies in the intersection of GL_1 with $\text{EL}_d(\bar{R})$, therefore the $1, 1$ entry is an invertible element in $\bar{R}$. $\square$

Proof of Lemma 1.2: Let $R = \text{Mat}_t(R')$ where R' is a finite commutative ring. Start with an element $g \in \text{EL}_d(R)$. Let $q = \lfloor d/2 \rfloor$ and $p = d - q \geq q$. Using that the ring $\text{Mat}_q(R)$ satisfies the Bass stable range condition we can find GEM

$$u = \begin{pmatrix} \text{Id}_q & * \\ 0 & \text{Id}_p \end{pmatrix}$$

such that $q \times q$ block in the upper right corner of ug is (left) invertible. Then there exist a second GEM u' such that lower right $q \times q$ block of $u'ug$ is equal to Id_q . Using left and right multiplication by two GEMs u'' and v we kill the two off-diagonal blocks of $u''u'ugv$. Thus we have shown that using 4 multiplications by generalized elementary matrices we can transform g to a matrix of type

$$\begin{pmatrix} * & 0 \\ 0 & \text{Id}_q \end{pmatrix}$$

where $q = \lfloor d/2 \rfloor$. If we apply the same procedure to the matrix in the upper left corner we can transform g to

$$\begin{pmatrix} * & 0 \\ 0 & \text{Id}_{q'} \end{pmatrix},$$

where $q' = \lfloor d/2 \rfloor + \lceil \lfloor d/2 \rfloor / 2 \rceil > d/2$ using 8 GEMs.

It is easy to see that for any $h \in \text{EL}_s(R) = \text{SL}_{st}(R')$ we have the identities

$$\begin{pmatrix} 0 & h \\ -h^{-1} & 0 \end{pmatrix} = \begin{pmatrix} 1 & h \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -h^{-1} & 1 \end{pmatrix} \begin{pmatrix} 1 & h \\ 0 & 1 \end{pmatrix} \quad \text{and}$$

$$\begin{pmatrix} h & 0 \\ 0 & h^{-1} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 1 - h^{-1} & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ h - 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & -h^{-1} \\ 0 & 1 \end{pmatrix},$$

in the group $\text{EL}_{2s}(R) = \text{SL}_{2st}(R')$.

Multiplying the three identities which result from the ones above when we substitute h_1 , $-h_2^{-1}$ and $h_1^{-1}h_2^{-1}$ for h in this order gives us that we can write

any matrix

$$\begin{pmatrix} [h_1, h_2] & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 0 & h_1 \\ -h_1^{-1} & 0 \end{pmatrix} \begin{pmatrix} 0 & -h_2^{-1} \\ h_2 & 0 \end{pmatrix} \begin{pmatrix} h_1^{-1}h_2^{-1} & 0 \\ 0 & h_2h_1 \end{pmatrix}$$

as a product of 10 generalized elementary matrices in $\text{EL}_{2s}(R)$.

Finally to complete the proof of Lemma 1.2 we only need to prove:

Lemma 3.3 *Let R' be a finite commutative ring. If $s > 2$ then any element in $\text{SL}_s(R')$ can be written as a product of 3 commutators.*

Without loss of generality we may assume that R' is a finite local commutative ring. In the case when R' is a field this is a result due to R. C. Thompson (see [6] for a survey of the related Ore Conjecture).

Theorem 3.4 *If K is a finite field and $\text{SL}_s(K)$ is perfect (i.e. $s > 2$ or $|K| > 3$) then every element of it is a commutator.*

Recall that by a result of Steinberg [15] the groups $\text{SL}_s(K)$ above can be generated by 2 elements. We shall need the following basic result:

Lemma 3.5 *Suppose that Γ is a group generated by $g_1, \dots, g_n$, and that V is a right $\mathbb{Z}\Gamma$ -module, such that $V = V(1 - \Gamma)$. Then in fact*

$$V = \sum_{i=1}^n V(1 - g_i).$$

Proof: Use that $1 - gg_i = g(1 - g_i) + (1 - g)$ in $\mathbb{Z}\Gamma$ and $g_i, g \in \Gamma$. $\square$

Proof of Lemma 3.3: Let I be the maximal ideal of R' , $K = R'/I$ and define $H_n := \ker(\pi_n : \text{SL}_s(R') \rightarrow \text{SL}_s(R'/I^n))$ with $H_0 = \text{SL}_s(R')$. It follows that $\bar{H} := H_0/H_1 \simeq \text{SL}_s(K)$ and by [15] it is generated by the images of some $a, b \in H_0$.

On the other hand, each of the quotients $V_n := H_n/H_{n+1}$ is an abelian group. We have that

$$[H_1, H_n] \leq H_{n+1} \quad \text{and} \quad [H_0, H_n] = H_n.$$

The first of these gives that the conjugation action of H_0 on V_n makes it into a right $\mathbb{Z}\bar{H}$ -module. The second implies that this module is *perfect*, i.e. that $V_n = V_n(\bar{H} - 1)$ for all n .

Claim: Every $g \in H_1$ can be written as

$$g = [a, x][b, y]$$

for appropriate $x, y \in H_1$.

Assuming this for the moment, Theorem 3.4 gives that any element $h \in \text{SL}_s(R')$ can be written as $h = [h_1, h_2]g$ with $g \in H_1$ and thus Lemma 3.3 follows immediately.

Proof of the claim: As $H_n = \{1\}$ for large enough n it is enough to prove the following:

Proposition 3.6 *Let $g \in H_1$ and $n \in \mathbb{N}$. There exist elements $x_n, y_n \in H_1$ such that*

$$g = [a, x_n][b, y_n] \mod H_n.$$

We shall prove this Proposition by induction starting with the case $n = 1$, where it is trivial. Assuming it is proved for some $n = m - 1$ we now prove it for $n = m$:

Without loss of generality, assume that $H_{m+1} = 1$, so that $V_m = H_m/H_{m+1}$ is central in H_1 .

Suppose $g = [a, x_n][b, y_n]g'$ for some $g' \in V_m$. We look for $x_{n+1} = \alpha x_n$, $y_{n+1} = \beta y_n$ with $\alpha, \beta \in V_m$.

Then by the commutator identities we have

$$\begin{aligned} [a, x_{n+1}][b, y_{n+1}] &= [a, \alpha x_n][b, \beta y_n] = [a, x_n][a, \alpha]^{x_n}[b, y_n][b, \beta]^{y_n} = \\ &= [a, x_n][b, y_n][a, \alpha][b, \beta], \end{aligned}$$

the last one because $\alpha, \beta, [a, \alpha], [b, \beta]$ are central in H_1 .

Therefore we only have to find $\alpha, \beta \in V_m$ such that

$$g' = \alpha(1 - a) + \beta(1 - b)$$

in the $\bar{H}$ -module V_m . The existence of such α and β is a consequence of Lemma 3.5, recalling that the images of a and $b \pmod{H_1}$ are generators for $\bar{H}$.

Proposition 3.6 is now proved, and thus Lemma 3.3 follows. $\square$

Let us finish the proof of Lemma 1.2. We have shown that any element g can be transformed to an element in $g' \in \text{EL}_{d'} = \text{SL}_{d't}(R')$ using 8 GEMs. If $d't > 2$ we can apply Lemma 3.3 to g' and write it as a product of 3 commutators, which can be expressed as a product of 10 GEMs each. This shows that we can write g as a product of $8 + 3 \cdot 10 = 38$ GEMs. If $d't \leq 2$ this argument does not work because we can not find a copy of $\text{SL}_6(R')$ embedded into $\text{EL}_d(R)$. If R is commutative we can apply Lemma 3.2 and see that every element in $\text{EL}_d(R)$ is a product of less than 20 generalized elementary matrices, because after the reduction we end up with the identity since $\text{GL}_1(R) \cap \text{EL}_d(R) = \{1\}$. Finally if $d = 3$ and $R = \text{Mat}_2(R')$ we need to modify the above argument slightly because the group $\text{SL}_2(\mathbb{F}_2)$ is not perfect. $\square$

4 Applications

4.1 Unbounded rank expanders

The matrix algebra $\text{Mat}_l(\mathbb{Z}[x_1, \dots, x_{sl}])$ can be generated as a ring by $s + 2$ elements for example the matrices

$$A = \sum_{i=1}^{l-1} e_{i,i+1} + (-1)^{l-1} e_{l,1} \quad B = e_{1,2} \quad Y_p = \sum_i x_{(p-1)l+i} e_{i,i}, \text{ for } p = 1, \dots, s.$$

Therefore we have a surjective homomorphism⁶

$$\mathbb{Z}\langle x_1, \dots, x_{s+2} \rangle \rightarrow \text{Mat}_l(\mathbb{Z}[x_1, \dots, x_{sl}]).$$

This gives a homomorphism

$$\pi_{l,s} : \text{EL}_3(\mathbb{Z}\langle x_1, \dots, x_{s+2} \rangle) \rightarrow \text{EL}_3(\text{Mat}_l(\mathbb{Z}[x_1, \dots, x_{sl}])) = \text{SL}_{3l}(\mathbb{Z}[x_1, \dots, x_{sl}]).$$

Let $S_{3,s+2}$ be the standard generating set of $G_{3,s+2}$, consisting of the 12 elementary matrices with ± 1 of the diagonal together with $8(s+2)$ elementary matrices with coefficients the generators $\pm x_i$ next to the main diagonal. Let $S'_{3l,sl}$ be the generating set of $\text{SL}_{3l}(\mathbb{Z}[x_1, \dots, x_{sl}])$ obtained by applying $\pi_{l,s}$ to $S_{3,s+2}$.

Proof of Theorem 11: Let $\rho : G = \text{SL}_{3l}(\mathbb{Z}[x_1, \dots, x_{sl}]) \rightarrow U(\mathcal{H})$ be a finite unitary representation and let v be ϵ -almost invariant vector for the set $S'_{3l,sl}$. Let $H = \ker \rho$ be its kernel. As in the proof of Theorem 6, let U be the ideal in $\mathbb{Z}[x_1, \dots, x_{sl}]$ generated by the off diagonal coefficients of the matrices in H .

Let $SK_1(R, U; 3l) = G(U)/\text{EL}(3l; U)$ where $G(U)$ is the principal congruence subgroup of G modulo U , i.e., the matrices in G congruent to Id_{3l} modulo U . We have the following diagram

$$\begin{array}{c} 1 \rightarrow SK_1(R, U; 3l) \rightarrow G/\text{EL}(3l; U) \rightarrow G(R/U) = \text{SL}_{3l}(\bar{R}) \rightarrow 1 \\ \downarrow \\ G/H \\ \downarrow \\ 1 \end{array}$$

where the row and the column are exact.

We can think of $\text{SL}_{3l}(\bar{R})$ as $\text{EL}_3(\text{Mat}_l(\bar{R}))$ and by Lemma 1.2 every element in this group can be written as a product of 38 generalized elementary matrices in $\text{EL}_3(\mathbb{Z}\langle x_1, \dots, x_{s+2} \rangle)$.

As in the proof of Theorem 6 every element in $SK_1(R, U; 3l)$ can be written as a product of $sl + 2$ Steinberg symbols which, as in the proof of Theorem 6, can be written as a product of $13(s+2)$ generalized elementary matrices.

Therefore every element in G can be written as a product of $64 + 13s$ GEMs and an element h in the kernel of the representation ρ . If we apply Lemma 1.1 to the group $\text{EL}_3(\mathbb{Z}\langle x_1, \dots, x_{s+2} \rangle)$ we can see that each of these GEMs moves the vector v by at most $2M(s+2)\epsilon$. If ϵ is small enough than every element in the group G moves v by less than $\sqrt{2}$, and the representation ρ has an invariant vector. This gives us a lower bound for the τ -constant of the form

$$\tau(\text{SL}_{3l}(\mathbb{Z}[x_1, \dots, x_{sl}]); S'_{3l,sl}) \geq \frac{1}{\sqrt{2}(64 + 13s)M(s+2)} \geq \frac{1}{400(4 + s^{3/2})},$$

which completes the proof of Theorem 11. $\square$

⁶There is also a surjective homomorphism $\mathbb{Z}\langle x_1, \dots, x_{s+2} \rangle \rightarrow \text{Mat}_l(\mathbb{Z}[x_1, \dots, x_{sl^2}])$, but we are unable to use it to obtain estimates of the Kazhdan constant for G_{l,sl^2} which do not depend on l .

Proof of Theorem 9: In the above construction if we put $s = 0$ we obtain the τ -constants for the groups $\mathrm{SL}_{3l}(\mathbb{Z})$ with respect to the generating sets $S'_{3l} = S'_{3l,0}$, consisting of 28 elements, are bounded below by $1/1600$.

If we are only interested in their finite images of the form $\mathrm{SL}_{3l}(\mathbb{F}_p)$ for $l > 3$ then τ -constant, which is equal to the Kazhdan constant because the group is finite, can be significantly improved. Let $g \in \mathrm{SL}_{3l}(\mathbb{F}_p) = \mathrm{SL}_3(\mathrm{Mat}_l(\mathbb{F}_p))$. Using 3 left multiplications by GEMs we can transform g to a 3×3 block matrix where the last column is trivial, with an extra 3 left multiplications by GEMs we can make the second column trivial. Finally with one GEM we can transform g to a block diagonal matrix where only the first entry is not trivial. However this entry is an element in $\mathrm{SL}_l(\mathbb{F}_p)$ and therefore is a commutator⁷ which is expressible as a product of 10 GEMs, i.e. we have:

$$\begin{aligned} \begin{pmatrix} * & * & * \\ * & * & * \\ * & * & * \end{pmatrix} &\xrightarrow{3} \begin{pmatrix} * & * & 0 \\ * & * & 0 \\ * & * & \mathrm{Id} \end{pmatrix} \xrightarrow{3} \begin{pmatrix} * & 0 & 0 \\ * & \mathrm{Id} & 0 \\ * & 0 & \mathrm{Id} \end{pmatrix} \xrightarrow{1} \\ &\xrightarrow{1} \begin{pmatrix} * & 0 & 0 \\ 0 & \mathrm{Id} & 0 \\ 0 & 0 & \mathrm{Id} \end{pmatrix} \xrightarrow{10} \begin{pmatrix} \mathrm{Id} & 0 & 0 \\ 0 & \mathrm{Id} & 0 \\ 0 & 0 & \mathrm{Id} \end{pmatrix} \end{aligned}$$

This shows that every matrix in $\mathrm{SL}_{3l}(\mathbb{F}_p)$ can be written as a product of 17 generalized elementary matrices coming from $\mathrm{SL}_3(\mathrm{Mat}_l(\mathbb{F}_p))$. Therefore if v is ϵ -invariant vector for the set S'_{3l} then g moves v by at most $17 \times 2M(2)\epsilon$, i.e., we have

$$\tau(\mathrm{SL}_{3l}(\mathbb{F}_p), S'_{3l}) = \mathcal{K}(\mathrm{SL}_{3l}(\mathbb{F}_p), S'_{3l}) \geq \frac{\sqrt{2}}{17 \times 2M(2)} \geq \frac{1}{400},$$

which completes the proof of Theorem 9.

4.2 Lubotzky-Wiess Conjectures

In [12] Lubotzky and Wiess conjectured that for a family of groups the property that their Cayley graphs are expanders is independent of the generating set. Recently in [1], Alon, Lubotzky and Wigderson disproved this conjecture using zig-zag products of graphs. They showed that the Cayley graphs of $\Gamma_p = \mathrm{SL}_2(\mathbb{F}_p) \ltimes F_2^{p+1}$ are not expanders with respect to some natural generating set, but that there are expanders with respect to a ‘random’ generating set. The proof is based on probabilistic arguments and does not give explicit generating sets which make the Cayley graphs expanders.

Theorem 9 together with the observation that the Cayley graphs of $\mathrm{SL}_{3l}(\mathbb{F}_p)$ are not expanders with respect to some natural generating sets, provides a natural and explicit counter example of Conjecture 7.

⁷This is not true if $p = 2$ and $l = 2$ because $\mathrm{SL}_2(\mathbb{F}_2)$ is not a perfect group. However it is easy to see that any element in this group is a product of 10 GEMs in $\mathrm{EL}_3(\mathrm{Mat}_2(\mathbb{F}_2))$.

In [12], Lubotzky and Wiess also conjectured that an infinite compact group K can not contain a finitely generated amenable dense subgroup and a f.g. dense subgroup having property T . This conjecture generalizes the observation that a discrete amenable group with property T is finite. The main motivation for this conjecture is the products of the groups $\mathrm{SL}_n(\mathbb{F}_p)$. It is known that in

$$K_n := \prod_p \mathrm{SL}_n(\mathbb{F}_p)$$

some natural f.g. dense subgroups (for example $\mathrm{SL}_n(\mathbb{Z})$) have property T and there is no f.g. dense amenable subgroup. On the other hand in

$$K^p := \prod_n \mathrm{SL}_n(\mathbb{F}_p)$$

some natural f.g. dense subgroups are amenable and there were no known subgroups with property T or τ .

Using non commutative universal lattices, we were able to construct a residually finite⁸ f.g. subgroup of K^p which have property τ . We believe that this group also has property T but we are unable to prove it. For technical reasons, we are going to work not with subgroups of K^p but with a similar infinite product.

Let $\widetilde{K}^p$ denote the infinite product

$$\widetilde{K}^p := \prod_{l \geq 4} \mathrm{SL}_{3l}(\mathbb{F}_p).$$

This is an infinite compact group, because it is an infinite product of finite, therefore compact, groups.

Let Γ_1 be the subgroup of $\widetilde{K}^p$ generated by A and B , where the components of A in each $\mathrm{SL}_{3l}(\mathbb{F}_p)$ are the matrices $A_{3l} = \sum e_{i,i+1} \pm e_{3l,1}$ and the components of B are $B_{3l} = \mathrm{Id}_{3l} + e_{1,2}$. Lubotzky and Wiess have shown:

Lemma 4.1 ([12], Example 4.3) *The group Γ_1 is finitely generated dense amenable subgroup of $\widetilde{K}^p$.*

Proof: First we will show that Γ_1 contains the infinite direct sum

$$\bigoplus_{l \geq 3} \mathrm{SL}_{3l}(\mathbb{F}_p) \subset \widetilde{K}^p,$$

which implies that Γ_1 is dense in $\widetilde{K}^p$.

It is enough to show that for each $l \geq 3$, the group Γ_1 contains the embedding of $\mathrm{SL}_{3l}(\mathbb{F}_p)$ in $\widetilde{K}^p$ – the proof uses induction on l . Suppose that we have shown this for all $l < k$. The direct computation shows that the group commutator of B and $A^{-n}BA^n$ has a nontrivial component in $\mathrm{SL}_{3s}(\mathbb{F}_p)$ if and only if $3s \mid n-1$ or

⁸Note that a discrete residually finite amenable group with property τ is also finite.

$3s \mid n+1$. Using the induction hypotheses and the element $[B, A^{-3k+1}BA^{3k-1}]$ we can see that Γ_1 contains element whose only non trivial component lies in the $\mathrm{SL}_{3k}(\mathbb{F}_p)$. The group $\mathrm{SL}_{3k}(\mathbb{F}_p)$ is quasi-simple and it is normally generated by any of its elements. This together with the observation that that Γ_1 projects onto $\mathrm{SL}_{3k}(\mathbb{F}_p)$ completes the induction step.

Let Δ_k be the subgroup of Γ_1 generated by $B, A^{-1}BA, \dots, A^{-k}BA^k$. It is easy to see that Δ_k is a finite group, thus $\Delta = \bigcup_k \Delta_k$ is an increasing union of finite groups and therefore is amenable. The same argument gives that the group $\tilde{\Delta} = \bigcup_i A^i \Delta A^{-i}$ is also amenable. By construction $\tilde{\Delta}$ is the normal closure of B in Γ_1 . Thus we have the exact sequence

$$1 \longrightarrow \tilde{\Delta} \longrightarrow \Gamma_1 \longrightarrow \mathbb{Z} \longrightarrow 1,$$

i.e., Γ_1 is extension of an amenable group by an amenable group and it is also amenable. $\square$

There is another way to look at the group $\widetilde{K^p}$

$$\widetilde{K^p} := \prod_{l \geq 2} \mathrm{SL}_{3l}(\mathbb{F}_p) = \prod_{l \geq 2} \mathrm{EL}_3(\mathrm{Mat}_l(\mathbb{F}_p)) = \mathrm{EL}_3 \left(\prod_{l \geq 2} \mathrm{Mat}_l(\mathbb{F}_p) \right).$$

The ring $R = \prod \mathrm{Mat}_l(\mathbb{F}_p)$ contains finitely generated dense subring S , thus group $\Gamma_2 = \mathrm{EL}_3(S)$ is a finitely generated dense subgroup of $\widetilde{K^p}$. The dense subring S is ‘almost’ isomorphic to the infinite direct sum

$$\bigoplus \mathrm{Mat}_l(\mathbb{F}_p)$$

and therefore any a finite quotient of $\mathrm{EL}_3(S)$ is ‘almost’ of the form $\mathrm{EL}_3(S')$ for some finite image S' of the ring S . This gives us that the finite images of Γ_2 are ‘almost’ finite product of the groups $\mathrm{SL}_{3l}(\mathbb{F}_p)$ for different l .

In the previous section we have shown that the Kazhdan constants, of these groups with respect to generating sets comming form a generating set of $\mathrm{EL}_3(S)$, are independent on l . This implies that the group $\mathrm{EL}_3(S)$ has property τ . There are several technical details, which need to be taken care of, in order to turn this idea into a proof.

Let S be the subring of $R = \prod \mathrm{Mat}_l(\mathbb{F}_p)$ generated by the elements $A, \bar{A}, C$, and D , where the components of A in each matrix algebra $\mathrm{Mat}_l(\mathbb{F}_p)$ are equal to $A_l = \sum e_{i,i+1} \pm e_{l,1}$, the components of $\bar{A}$ are A_l^{-1} ; the components of C and D are $C_l = e_{1,2}$ and $D_l = e_{2,1}$ respectively.

Lemma 4.2 *The ring S contains the infinite direct sum*

$$\bigoplus_{l \geq 3} \mathrm{Mat}_l(\mathbb{F}_p)$$

and it is a dense subring of R .

Proof: It is enough to show that for any l the ring S contains the embedding of the matrix algebra $\text{Mat}_l(\mathbb{F}_p)$ in R – the proof uses induction on l . Suppose that we have shown this for all $l < k$. The direct computation shows that the ring commutator of C and $\bar{A}^n C A^n$ has nontrivial component in $\text{Mat}_s(\mathbb{F}_p)$ if and only if $s \mid n - 1$ or $s \mid n + 1$. Using the induction hypotheses and the element $[C, \bar{A}^{k-1} C A^{k-1}]$ we can see that S contains an element whose only non trivial component lies in the $\text{Mat}_k(\mathbb{F}_p)$. The algebra $\text{Mat}_k(\mathbb{F}_p)$ is simple and the two-sided ideal generated by any non-zero element coincides with the whole algebra. This together with the observation that that S projects onto $\text{Mat}_k(\mathbb{F}_p)$ completes the induction step. $\square$

Let Γ_2 be the subgroup $\text{EL}_3(S) \subset \widetilde{K}^p = \text{EL}_3(R)$. This group is dense because S is a dense subring of R . We also have that S is a factor ring of $\mathbb{Z}\langle x_1, x_2, x_3, x_4 \rangle$, thus Γ_2 is generated by the image of the set $S_{3,4}$, i.e., by the matrices

$$e_{i,j}(\pm 1), \quad e_{i,j}(\pm A), \quad e_{i,j}(\pm \bar{A}), \quad e_{i,j}(\pm C) \quad \text{and} \quad e_{i,j}(\pm D).$$

This gives us a finitely generated dense subgroup in $\widetilde{K}^p$. Theorem 10 will follow from Lemmas 4.1 and 4.2 and:

Lemma 4.3 *The group Γ_2 has property τ .*

Proof: Let ρ be a finite representation of Γ_2 and let $H = \ker \rho$. As in the proof of Theorem 6 let U be the subset:

$$U = \{u \in S \mid e_{i,j}(u) \in H, \text{ for some } i \neq j\}.$$

This is a two sided ideal of finite index in S . By construction A is invertible element in the ring S , therefore there exists N such that $A^N - 1 \in U$. Let I_N be the two-sided ideal in S generated by $A^N - 1$. The inclusion $I_N \subset U$ leads us to the diagram:

$$\begin{array}{c} 1 \rightarrow SK_1(S, I_N; 3) \rightarrow \Gamma_2 / \text{EL}_3(I_N) \rightarrow \text{EL}_3(S/I_N) \rightarrow 1. \\ \downarrow \\ \Gamma_2 / H \end{array}$$

In order to finish the proof that Γ_2 has property τ it is sufficient to prove that the groups $\text{EL}_3(S/I_N)$, for all N , have uniform bounded elementary generation property, and that any element in the K -group $SK_1(S, I_N; 3)$ can be written as a product of small number of elementary matrices.

Let us consider the factor ring S/I_N :

Lemma 4.4 *The factor ring S/I_N is isomorphic to*

$$S/I_N \simeq \mathbb{F}_p[C_N] \oplus \bigoplus_{l \mid N, l \geq 3} \text{Mat}_l(\mathbb{F}_p),$$

where $\mathbb{F}_p[C_N]$ is the group algebra of the cyclic group C_N .

Proof: The inclusion $\bigoplus \text{Mat}_l(\mathbb{F}_p) \subset S$ implies that

$$S \simeq S^{>2N} \oplus \bigoplus_{l \geq 3}^{2N} \text{Mat}_l(\mathbb{F}_p)$$

where $S^{>2N}$ is the the image of S in $\prod_{l > 2N} \text{Mat}_l \mathbb{F}_p$. This isomorphism implies that

$$\begin{aligned} S/I_N &\simeq S^{>2N} / (I_N \cap S^{>2N}) \oplus \bigoplus_{l \geq 3}^{2N} \text{Mat}_l(\mathbb{F}_p) / (I_N \cap \text{Mat}_l(\mathbb{F}_p)) \\ &\simeq S^{>2N} / I_N^{>2N} \oplus \bigoplus_{l \geq 3}^{2N} \text{Mat}_l(\mathbb{F}_p) / I_N^l, \end{aligned}$$

where I_N^l is the two-sided ideal $\text{Mat}_l(\mathbb{F}_p)$ in generated by $A_l^N - 1$ and $I_N^{>2N}$ is the two-sided ideal in $S^{>2N}$ generated by the image of $A^N - 1$.

The algebra $\text{Mat}_l(\mathbb{F}_p)$ does not have any nontrivial ideals, i.e., if $A_l^N - 1 \neq 0$ then $I_N^l = \text{Mat}_l(\mathbb{F}_p)$. The last condition is equivalent to $l|N$. Therefore we have the isomorphism:

$$\bigoplus_{l \geq 3}^{2N} \text{Mat}_l(\mathbb{F}_p) / I_N^l \simeq \bigoplus_{l|N, l \geq 2} \text{Mat}_l(\mathbb{F}_p).$$

Using the definition of A , $\bar{A}$, C and D it can be verified that in the ring $S^{>2N}$ we have the identities:

$$\begin{aligned} [C, \bar{A}CA], \bar{A}DA] &= C, & [C, \bar{A}^{N+1}CA^{N+1}] &= 0, \\ [\bar{A}DA, D], \bar{A}CA] &= D, & [D, \bar{A}^{N+1}DA^{N+1}] &= 0, \end{aligned}$$

because these identities hold in $\text{Mat}_l(\mathbb{F}_p)$ for all $l > 2N$. These identities imply that

$$\begin{aligned} C &\equiv [[C, \bar{A}CA], \bar{A}DA] \equiv [[C, \bar{A}^{N+1}CA^{N+1}], \bar{A}DA] \equiv 0 \pmod{I_N^{>2N}} \\ D &\equiv [[\bar{A}DA, D], \bar{A}CA] \equiv [[\bar{A}^{N+1}DA^{N+1}, D], \bar{A}CA] \equiv 0 \pmod{I_N^{>2N}}, \end{aligned}$$

i.e., $S^{>2N} / I_N^{>2N}$ is generated by the images of A and $\bar{A}$. Therefore the factor ring is isomorphic to

$$\mathbb{Z}\langle A, \bar{A} \rangle / (A.\bar{A} = 1, A^N = 1, p = 0).$$

The last ring is isomorphic to the group algebra $\mathbb{F}_p[C_N]$. $\square$

Therefore we have

$$\text{EL}_3(S/I_N) = \text{EL}_3(\mathbb{F}_p[C_N]) \times \prod_{l|N} \text{EL}_3(\text{Mat}_l(\mathbb{F}_p)).$$

By Lemma 1.2 any element in each term is a product of 40 elementary matrices, i.e., we have that $BE_3(S/I_N) < 40$.

It remains to prove the group $SK_1(S, I_N; 3)$ is small. Using K -theory we can see that this is an image of $K_2(S/I_N)$. The basic properties of the functor K_2 yield

$$K_2(S/I_N) = K_2(\mathbb{F}_p[C_N]) \oplus \bigoplus_{l|N} K_2(\text{Mat}_l(\mathbb{F}_p)).$$

It is known that $K_2(\text{Mat}_l(\mathbb{F}_p)) = K_2(\mathbb{F}_p) = 0$, therefore

$$K_2(S/I_N) = K_2(\mathbb{F}_p[C_N]).$$

The ring $\mathbb{F}_p[C_N]$ is a finite commutative ring, generated by 1 and another element. Lemma 1.3 gives that any element in $K_2(\mathbb{F}_p[C_N])$ is a product of 3 Steinberg symbols⁹ and can be written as a product of 39 elementary matrices.

This shows any element of Γ_2 can be written as a product of 80 elementary matrices modulo an element in the kernel of ρ . Applying Theorem 5 gives us that Γ_2 has property τ and its τ -constant is at least

$$\tau(\Gamma_2, \{e_{i,j}(\pm 1), e_{i,j}(\pm A), e_{i,j}(\pm \bar{A}), e_{i,j}(\pm C), e_{i,j}(\pm D) \mid 1 \leq i \neq j \leq 3\}) > \frac{1}{2200}.$$

This completes the proof of Lemma 4.3. $\square$

As we described above Theorem 10 follows immediately from Lemmas 4.1, 4.2 and 4.3.

References

- [1] N. Alon, A. Lubotzky, A. Wigderson, Semi-direct product in groups and zig-zag product in graphs: connections and applications, *42nd IEEE Symposium on Foundations of Computer Science (Las Vegas, NV, 2001)*, 630–637, *IEEE Computer Soc., Los Alamitos, CA, 2001*.
- [2] M. Burger, Kazhdan constants for $SL_3(\mathbb{Z})$, *J. Reine Angew. Math.*, 413 (1991), 36–67.
- [3] D. Carter, G. Keller, Bounded elementary generation of $SL_n(\mathcal{O})$. *Amer. J. Math* 105 (1983), no.3, 673–687.
- [4] R. K. Dennis, M. R. Stein, K_2 of discrete valuation rings. *Advances in Math.* 18 (1975), no. 2, 182–238.
- [5] R. K. Dennis, L. N. Vaserstein, On a question of M. Newman on the number of commutators. *J. Algebra* 118 (1988), no. 1, 150–161.
- [6] E. W. Ellers, N. Gordeev, On the conjectures of J. Thompson and O. Ore, *Trans. Amer. Math. Soc.* 350 (1998), 3657–3671.

⁹Using the results in [4], it can be shown that $K_2(\mathbb{F}_p[C_N])$ is a trivial group

- [7] M. Kassabov, Kazhdan constants for $SL_d(\mathbb{Z})$, [arXiv:math.GR/0311487](#), to appear in *Internat. J. Algebra and Comput.*
- [8] M. Kassabov, N. Nikolov, Universal lattices and property τ , [arXiv:math.GR/0502112](#).
- [9] M. Kassabov, T. R. Riley, Diameter of Cayley graphs of $SL_n(\mathbb{Z}/k\mathbb{Z})$, [arXiv:math.GR/0502221](#).
- [10] D. A. Kazhdan, On the connection of the dual space of a group with the structure of its closed subgroups, *Funkcional. Anal. i Priloz. 1 (1967)*, 71-74.
- [11] A. Lubotzky, *Discrete groups, expanding graphs and invariant measures*, Progr. Math. 125, Birkhäuser, Boston, 1994.
- [12] A. Lubotzky, B. Weiss, Groups and expanders, *Expanding graphs, DIMACS Ser. Discrete Math. Theoret. Comput. Sci., 10, Amer. Math. Soc., Providence, RI, 1993*.
- [13] A. Lubotzky, A. Żuk, *On Property (τ)* , book.
- [14] Y. Shalom, Bounded generation and Kazhdan's property (T), *Publ. Math. IHES, 90 (1999)*, 145-168.
- [15] R. P. Steinberg, Generators for finite simple groups, *Canad. J. Math. 14 (1962)*, 277-283.
- [16] L. N. Vaserstein, On the stabilization of the general linear group over a ring. *Mat. Sb. (N.S.) 79 (121) 405-424 (Russian); translated as Math. USSR-Sb 8 1969 383-400*.
- [17] L. N. Vaserstein, Bass's first stable range condition. *J. Pure Appl. Algebra 34:2-3 (1984) 319-330*.

Martin Kassabov,
 Cornell University, Ithaca, NY 14853-4201, USA.
e-mail: kassabov@math.cornell.edu